



(51) International Patent Classification <sup>6</sup> : <b>A61K 4900, C12N 7/00, 7/02, 15/00, 15/06</b>		<b>A1</b>	(11) International Publication Number: <b>WO 97/04814</b> (43) International Publication Date: 13 February 1997 (13.02.97)
(21) International Application Number: PCT/US96/12510 (22) International Filing Date: 30 July 1996 (30.07.96) (30) Priority Data: 08/509,261 31 July 1995 (31.07.95) US 08/521,992 31 August 1995 (31.08.95) US (71) Applicant: THE REGENTS OF THE UNIVERSITY OF CALIFORNIA (US/US); Office of Technology Transfer, Suite 150, 1320 Harbor Bay Parkway, Alameda, CA 94501 (US). (72) Inventors: PRUSINER, Stanley, B.; 400 Pacheco Street, San Francisco, CA 94116 (US). SCOTT, Michael, R.; 1200 Clayton Street #9, San Francisco, CA 94114 (US). TELLING, Glenn; 2140 Fell Street #302, San Francisco, CA 94117 (US). (74) Agent: BOZICEVIC, Karl; Fish & Richardson P.C., Suite 100, 2200 Sand Hill Road, Menlo Park, CA 94025 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  Published With international search report.	
(54) Title: DETECTING PRIONS IN A SAMPLE AND PRION PREPARATION AND TRANSGENIC ANIMAL USED FOR SAME			
(57) Abstract  The invention includes an artificial PrP gene, a transgenic animal containing a Prp gene of a heterologous animal, and a hybrid non-human animal with an ablated endogenous prion protein gene. Also disclosed are methods of assaying for the presence of prion particles in a sample by inoculating the claimed transgenic animals and observing the inoculated animals for symptoms associated with prion diseases.			

- 1 -

DETECTING PRIONS IN A SAMPLE AND PRION  
PREPARATION AND TRANSGENIC ANIMAL USED FOR SAME

5

Government Rights

The United States Government may have certain rights in this application pursuant to Grant Nos. NS14069, AG02132, NS22786, AG08967 and AG10770 awarded by the National Institutes of Health.

10

Field of the Invention

This invention relates generally to chimeric genes, methods of assaying and to transgenic animals used in such assays. More specifically, this invention relates to artificial and chimeric PrP genes, assaying samples for pathogenic prions, standardized prion preparations used in such assays and to transgenic mice and hybrid transgenic mice which can be infected with prions which generally only infect a genetically diverse species.

15  
20

Background of the Invention

Prions are infectious pathogens that cause central nervous system spongiform encephalopathies in humans and animals. Prions are distinct from bacteria, viruses and viroids. The predominant hypothesis at present is that

94

no nucleic acid component is necessary for infectivity of prion protein. Further, a prion which infects one species of animal (e.g., a human) will not infect another (e.g., a mouse).

25  
30

A major step in the study of prions and the diseases that they cause was the discovery and purification of a protein designated prion protein ("PrP") [Bolton et al. Science 218:1309-11 (1982); Prusiner et al., Biochemistry 21:6942-50 (1982); McKinley

35

SUBSTITUTE SHEET (RULE 26)

- 2 -

et al., Cell 35:57-62 (1983)]. Complete prion protein-encoding genes have since been cloned, sequenced and expressed in transgenic animals. PrP<sup>C</sup> is encoded by a single-copy host gene [Basler et al., Cell 46:417-28  
5 (1986)] and is normally found at the outer surface of neurons. A leading hypothesis is that prion diseases result from conversion of PrP<sup>C</sup> into a modified form called PrP<sup>Sc</sup>. However, the actual biological or physiological function of PrP<sup>C</sup> is not known.

10 It appears that the scrapie isoform of the prion protein (PrP<sup>Sc</sup>) is necessary for both the transmission and pathogenesis of the transmissible neurodegenerative diseases of animals and humans. See Prusiner, S.B., "Molecular biology of prion disease," Science 252:1515-  
15 1522 (1991). The most common prion diseases of animals are scrapie of sheep and goats and bovine spongiform encephalopathy (BSE) of cattle [Wilesmith, J. and Wells, Microbiol. Immunol. 172:21-38 (1991)]. Four prion diseases of humans have been identified: (1) kuru,  
20 (2) Creutzfeldt-Jakob Disease (CJD), (3) Gerstmann-Strassler-Scheinker Disease (GSS), and (4) fatal familial insomnia (FFI) [Gajdusek, D.C., Science 197:943-960 (1977); Medori et al., N. Engl. J. Med. 326:444-449 (1992)]. The presentation of human prion diseases as  
25 sporadic, genetic and infectious illnesses initially posed a conundrum which has been explained by the cellular genetic origin of PrP.

Most CJD cases are sporadic, but about 10-15% are inherited as autosomal dominant disorders that are caused  
30 by mutations in the human PrP gene [Hsiao et al., Neurology 40:1820-1827 (1990); Goldfarb et al., Science 258:806-808 (1992); Kitamoto et al., Proc. R. Soc. Lond. (In press) (1994)]. Iatrogenic CJD has been caused by human growth hormone derived from cadaveric pituitaries  
35 as well as dura mater grafts [Brown et al., Lancet

- 3 -

340:24-27 (1992)]. Despite numerous attempts to link CJD to an infectious source such as the consumption of scrapie infected sheep meat, none has been identified to date [Harries-Jones et al., J. Neurol. Neurosurg. Psychiatry 51:1113-1119 (1988)] except in cases of iatrogenically induced disease. On the other hand, kuru, which for many decades devastated the Fore and neighboring tribes of the New Guinea highlands, is believed to have been spread by infection during ritualistic cannibalism [Alpers, M.P., Slow Transmissible Diseases of the Nervous System, Vol. 1, S.B. Prusiner and W.J. Hadlow, eds. (New York: Academic Press), pp. 66-90 (1979)].

The initial transmission of CJD to experimental primates has a rich history beginning with William Hadlow's recognition of the similarity between kuru and scrapie. In 1959, Hadlow suggested that extracts prepared from patients dying of kuru be inoculated into non-human primates and that the animals be observed for disease that was predicted to occur after a prolonged incubation period [Hadlow, W.J., Lancet 2:289-290 (1959)]. Seven years later, Gajdusek, Gibbs and Alpers demonstrated the transmissibility of kuru to chimpanzees after incubation periods ranging from 18 to 21 months [Gajdusek et al., Nature 209:794-796 (1966)]. The similarity of the neuropathology of kuru with that of CJD [Klatzo et al., Lab Invest. 8:799-847 (1959)] prompted similar experiments with chimpanzees and transmissions of disease were reported in 1968 [Gibbs, Jr. et al., Science 161:388-389 (1968)]. Over the last 25 years, about 300 cases of CJD, kuru and GSS have been transmitted to a variety of apes and monkeys.

The expense, scarcity and often perceived inhumanity of such experiments have restricted this work and thus limited the accumulation of knowledge. While



the most reliable transmission data has been said to emanate from studies using non-human primates, some cases of human prion disease have been transmitted to rodents but apparently with less regularity [Gibbs, Jr. et al.,

5 Slow Transmissible Diseases of the Nervous System,

Vol. 2, S.B. Prusiner and W.J. Hadlow, eds. (New York: Academic Press), pp. 87-110 (1979); Tateishi et al., Prion Diseases of Humans and Animals, Prusiner et al., eds. (London: Ellis Horwood), pp. 129-134 (1992)].

- 10           The infrequent transmission of human prion disease to rodents has been cited as an example of the "species barrier" first described by Pattison in his studies of passaging the scrapie agent between sheep and rodents [Pattison, I.H., NINDB Monograph 2, D.C. Gajdusek,
- 15 C.J. Gibbs Jr. and M.P. Alpers, eds. (Washington, D.C.: U.S. Government Printing), pp. 249-257 (1965)]. In those investigations, the initial passage of prions from one species to another was associated with a prolonged incubation time with only a few animals developing
- 20 illness. Subsequent passage in the same species was characterized by all the animals becoming ill after greatly shortened incubation times.

          The molecular basis for the species barrier between Syrian hamster (SHa) and mouse was shown to

25 reside in the sequence of the PrP gene using transgenic (Tg) mice [Scott et al., Cell 59:847-857 (1989)]. SHaPrP differs from MoPrP at 16 positions out of 254 amino acid residues [Basler et al., Cell 46:417-428 (1986); Loch et al., Proc. Natl. Acad. Sci. USA 83:6372-6376 (1986)].

30 Tg(SHaPrP) mice expressing SHaPrP had abbreviated incubation times when inoculated with SHa prions. When similar studies were performed with mice expressing the human, or ovine PrP transgenes, the species barrier was not abrogated, i.e., the percentage of animals which

35 became infected were unacceptably low and the incubation

- 5 -

times were unacceptably long. Thus, it has not been possible, for example in the case of human prions, to use transgenic animals (such as mice containing a PrP gene of another species) to reliably test a sample to determine  
5 if that sample is infected with prions. The seriousness of the health risk resulting from the lack of such a test is exemplified below.

More than 45 young adults previously treated with HGH derived from human pituitaries have developed CJD  
10 [Koch et al., N. Engl. J. Med. 313:731-733 (1985); Brown et al., Lancet 340:24-27 (1992); Fradkin et al., JAMA 265:880-884 (1991); Buchanan et al., Br. Med. J. 302:824-828 (1991)]. Fortunately, recombinant HGH is now used, although the seemingly remote possibility has been raised  
15 that increased expression of wtPrP<sup>C</sup> stimulated by high HGH might induce prion disease [Lasmezas et al., Biochem. Biophys. Res. Commun. 196:1163-1169 (1993)]. That the HGH prepared from pituitaries was contaminated with prions is supported by the transmission of prion disease  
20 to a monkey 66 months after inoculation with a suspect lot of HGH [Gibbs, Jr. et al., N. Engl. J. Med. 328:358-359 (1993)]. The long incubation times associated with prion diseases will not reveal the full extent of iatrogenic CJD for decades in thousands of people treated  
25 with HGH worldwide. Iatrogenic CJD also appears to have developed in four infertile women treated with contaminated human pituitary-derived gonadotrophin hormone [Healy et al., Br. J. Med. 307:517-518 (1993); Cochius et al., Aust. N.Z. J. Med. 20:592-593 (1990);  
30 Cochius et al., J. Neurol. Neurosurg. Psychiatry 55:1094-1095 (1992)] as well as at least 11 patients receiving dura mater grafts [Nisbet et al., J. Am. Med. Assoc. 261:1118 (1989); Thadani et al., J. Neurosurg. 69:766-769 (1988); Willison et al., J. Neurosurg. Psychiatric 54:940  
35 (1991); Brown et al., Lancet 340:24-27 (1992)]. These

- 6 -

cases of iatrogenic CJD underscore the need for screening pharmaceuticals that might possibly be contaminated with prions.

Recently, two doctors in France were charged with  
5 involuntary manslaughter of a child who had been treated  
with growth hormones extracted from corpses. The child  
developed Creutzfeldt-Jakob Disease. (See New Scientist,  
July 31, 1993, page 4). According to the Pasteur  
Institute, since 1989 there have been 24 reported cases  
10 of CJD in young people who were treated with human growth  
hormone between 1983 and mid-1985. Fifteen of these  
children have died. It now appears as though hundreds of  
children in France have been treated with growth hormone  
extracted from dead bodies at the risk of developing CJD  
15 (see New Scientist, November 20, 1993, page 10.) In  
view of such, there clearly is a need for a convenient,  
cost-effective assay for testing sample materials for the  
presence of prions which cause CJD. The present  
invention offers such an assay.

## 20 Summary of the Invention

The invention comprises a standardized prion  
preparation, chimeric PrP genes with mutation codons,  
transgenic mice which can be used in preparing such a  
standardized preparation and methods of testing samples  
25 using the preparation and transgenic mice. In order to  
produce the standardized prion preparation it is  
necessary to produce a group of  
non-human host mammals which have their genome  
manipulated with respect to genetic material related to a  
30 PrP gene such that the mammals are susceptible to  
infection with a prion which generally only infects an  
animal which is genetically diverse from the host. The  
transgenic host animals produced are inoculated with a  
prion containing composition and the animals are observed

- 7 -

until they exhibit symptoms of prion infection. Brain tissue is harvested from the animals and homogenized to create the standardized prion preparation. This process can be repeated one more time using homogenized brain tissue of the last inoculated group to inoculate a new group and thereby further standardize the preparation and reduce any irregularities that might be created by the composition of the initial prion inoculation composition. Different forms of transgenic animals can be used in the production of different preparations and two or more different standardized preparation can be mixed.

However, it is preferable to produce the preparation using a non-human mammal which has its endogenous PrP gene ablated and includes an exogenous PrP gene which can be the PrP gene from a genetically diverse species, an artificial PrP gene which includes a portion of the PrP gene of a genetically diverse species or a completely artificial PrP gene. To test samples for the presence of prions two sets of non-human transgenic mammals are prepared. Both sets of mammals are designed so that they are susceptible to infection by prions which would normally only infect a genetically diverse species. The first set of animals are inoculated with a standard prion preparation and the second set is inoculated with the test sample. Both sets are observed and the set of mammals which is inoculated with a standard prion preparation is used as a control. If the group of animals inoculated with the test sample develop symptoms of prion infection then the tester can deduce that the sample includes prions. If the group inoculated with the test sample does not develop symptoms of prion disease and the group inoculated with the standard prion preparation does then the absence of prions in the sample is deduced.

- 8 -

The invention includes a chimeric artificial PrP gene comprising portions of the host animal e.g., end portions of a host animal and a middle portion of a genetically diverse test animal wherein the middle  
5 portion includes a specific alterations designed to match that of a disease state of such a host. Further, the invention includes, a transgenic animal containing the artificial gene or elevated expression of a PrP gene from a genetically diverse animal, hybrid transgenic animals  
10 which are the offspring of different transgenic animals with each other or with a transgenic animal that has an ablated endogenous prion protein gene, a standardized prion preparation and assay methodology which uses the preparation and genetically altered animals to detect  
15 pathogenic prions in a sample.

The artificial gene includes a sequence such that when it is inserted into the genome of an animal (such as a mouse), the animal is rendered susceptible to infection with prions which normally would infect only a specific  
20 species of genetically diverse animal (such as a human, cow, sheep, pig, chicken, cat or dog). The artificial PrP gene may be comprised partially or completely of an artificial polynucleotide sequence, i.e. codon sequences not present in any native PrP gene sequence.

25 Alternatively, the artificial gene may be comprised of the codon sequence of a host animal with one or more codon substitutions being made wherein the substitutions are preferably corresponding PrP gene codons from a genetically diverse animal, meaning that PrP gene differs  
30 from the PrP gene of the host animal by 20 or more codons. Transgenic animals containing elevated levels of expression of the PrP gene which can be obtained for example, by over expression of the gene with an enhanced promoter and/or with high copy numbers of the natural PrP  
35 gene of a genetically diverse animal are also disclosed.



Hybrid transgenic animals include animals resulting from a cross between two transgenic animals and in particular a cross between a transgenic animal containing the entire prion protein gene of a genetically diverse animal (e.g.,  
5 a mouse containing a human prion protein gene) and an animal with its endogenous prion protein gene disrupted (e.g., a mouse with an ablated prion protein gene). Hybrids also specifically include crossing a transgenic animal having a chimeric prion protein gene with an  
10 animal with its endogenous prion protein gene ablated.

Genetics constructs and methodologies of the invention are used to create animals which due to their genetic make up will develop disease from inoculation with prions which would generally only infect a  
15 genetically diverse animal, e.g., a mouse of the invention will consistently become infected with prions which generally will only infect a human and symptoms of the infection will become apparent in a short period e.g., 350 days or less. The animals of the invention are  
20 used in assays to test samples of any given material to determine if the material includes prions which would infect another animal (such as a human) if the material were ingested or injected. Standardized prion preparations of the invention are used to inoculate  
25 animals of the invention to create controls when carrying out an assay of the invention. The standardized prion preparation will always contain prions which will infect a genetically modified animal of the invention which animal will develop clinical signs of CNS dysfunction  
30 within a set period of time.

In one preferred example the mouse genome includes a chimeric PrP gene which gene includes a portion of a gene of the animal (e.g. human) in danger of infection from prions in the sample. For example, Creutzfeldt  
35 Jakob Disease (CJD) is a fatal neurodegenerative disease



- 10 -

of humans caused by prions. Preferred transgenic (Tg) mice disclosed herein express a chimeric prion protein (PrP) in which a segment of mouse (Mo) PrP was replaced with the corresponding human (Hu) PrP sequence. The  
5 chimeric PrP designated MHu2MPrP, differs from MoPrP by 9 codons between codons 96 and 167. All of the Tg(MHu2MPrP) mice injected with human prions developed neurologic disease. More specifically, the transgenic mice of the invention developed the disease ~200 days  
10 after inoculation with brain homogenates from three CJD patients. When inoculated with CJD prions, MHu2MPrP<sup>sc</sup> was formed; in contrast MoPrP<sup>sc</sup> was produced if Mo prions were inoculated. Tg(MHu2MPrP) mice disclosed herein are useful in the diagnosis, prevention and treatment of  
15 human prion diseases. Transgenic mice containing the artificial PrP gene or elevated levels of expression of a native PrP gene from a genetically diverse animal can be used to test samples for prions which might infect such animals. The transgenic and hybrid animals disclosed  
20 herein consistently develop the adverse effects of such prions in a relatively short time and are substantially cheaper and easier to maintain than are currently used primate models. Transgenic mice containing a human prion protein gene are designated Tg(HuPrP) and may be crossed  
25 with mice with an ablated endogenous prion protein gene which are designated Prnp<sup>0/0</sup> to obtain a hybrid designated Tg(HuPrP)/Prnp<sup>0/0</sup>.

An important object of the invention is to provide a standardized prion preparation which is produced by  
30 inoculating a non-human host animal which has its genome manipulated with respect to its PrP gene so that it is susceptible to infection with prions which generally only infect an animal genetically diverse from the host animal. The host animal is inoculated with prions and  
35 the animal observed until symptoms of infection occur

- 11 -

after which brain tissue is harvested from the animal and homogenized to produce the standardized prion preparation.

Another object of the invention is to provide a  
5 transgenic, hybrid, non-human mammal which has its endogenous PrP gene ablated and which includes a PrP gene from a genetically diverse mammal or a manipulated PrP gene such as a chimeric PrP gene comprised of codons from the host mammal and a genetically diverse mammal.

10 Yet another object of the invention is to provide for a method of testing samples for the presence of prions. The method involves creating two groups of non-human mammals which have their genome altered so that they are susceptible to infection with prions which  
15 generally only infect a genetically diverse animal. The first group of animals is infected with a test sample and the second group is infected with a standardized prion preparation. Both groups of mammals are observed in the presence of prions and the sample can be deduced if the  
20 first group of animals develop symptoms of prion infection.

An advantage of the invention is that a standardized prion preparation can be used to provide a control group when testing samples for the presence of  
25 prions.

Another object is to provide a hybrid animal which is obtained by crossing an animal having an ablated en-  
30 geneous prion protein gene with a transgenic animal containing (1) a chimeric gene or (2) the prion protein gene of a genetically diverse animal which gene may be present at elevated levels.

Another object is to provide a standardized prion preparation produced from harvested brain tissue taken from animals of the invention (that have substantially  
35 identical genomes and specifically have substantially

- 12 -

identical genetic material related to prions) which animals exhibit symptoms of prion infection after being inoculated with prions which generally only infect a genetically diverse species.

- 5           A feature of the invention is that the standardized prion preparations of the invention can be used to consistently inoculate control animals with a known amount and type of prion.

          A feature of the invention is that the PrP gene of  
10 the host animal can be altered by replacing codons with codons of a test animal at the same relative position which differ from the codons of the host animal, up to and including replacing all the differing codons wherein the codons are replaced in a manner so as to maintain the  
15 operability of the gene.

          Another object is to provide an artificial PrP gene wherein one or more codons (preferably 1-39 codons) of the PrP gene of a host animal (e.g. a mouse) is replaced with codons of the PrP gene of a genetically  
20 diverse test animal (e.g. a human, cow or sheep) in a manner so as to render the host animal susceptible to infection with prions which normally infect only the genetically diverse test animal.

          Another object is to provide a chimeric gene  
25 comprised of codons encoding the C- and N- terminus of one species of mammal and middle codons of another species of mammal.

          Another object of the invention is to provide a transgenic host mammal (which is small, e.g., less than  
30 1 kg when full grown, and inexpensive to maintain) such as a mouse, rat or hamster which includes a chimeric PrP gene which gene includes a portion of the PrP gene from another animal, (which is large, greater than 2 kg when full grown, and expensive to maintain) such as a human,  
35 cow, pig, sheep, cat or dog.

- 13 -

Another object of the invention is to provide a transgenic host animal which includes elevated levels of expression of a native PrP gene of a genetically diverse animal wherein the elevated levels of expression are  
5 obtained by the inclusion of a high copy number of the PrP gene of the genetically diverse mammal and/or fusing an enhanced promoter to the PrP gene of the genetically diverse animal which transgenic animal may be used by itself to assay for prions or for cross-breeding with an  
10 animal which has an ablated endogenous prion protein gene.

An advantage of the present invention is that the transgenic and hybrid animal can be used to assay for the presence of prions in a sample in a manner which is  
15 substantially faster, more efficient and cheaper than presently available assay methods.

Another advantage is that transgenic and hybrid animals inoculated with prions of humans can be used as test animals for testing drugs for efficacy in the  
20 treatment of humans suffering from diseases resulting from infection with prions.

Another advantage is that the transgenic and hybrid animals can detect prions in a sample at very low levels, e.g., 1 part per million, and even as low as  
25 1 part per billion.

Still another advantage is that the transgenic and hybrid animals provide an assay which is highly accurate, i.e., does not provide false positives and consistently determines the presence of prions.

30 Yet another advantage is that by increasing the copy number of an exogenous prion protein gene of the invention in a transgenic or hybrid and/or disrupting the endogenous gene of, the incubation time for prion caused disease is decreased.

- 14 -

Another advantage is that the standardized prion preparations of the invention can eliminate the need for extracting brain tissue from mammals which may have been infected with different types of prions and may each have a different genetic make up regarding genetic material related to prions.

Another advantage is that assays of the invention can be carried out more reliably using the standardized prion preparations of the invention.

10 A feature of the present invention is that the transgenic and hybrid animals injected with a sample containing pathogenic prions will consistently develop the disease effects of the prions within a relatively short time, e.g. about 200 days  $\pm$  50 days after injection  
15 or less.

Another feature is that an artificial gene of the invention preferably contains codons of the PrP gene of a host animal (such as a mouse) with some (but not all) of the codons which differ from the mouse and a second  
20 genetically diverse test mammal (such as a human) replacing codons of the first mammal at the same relative positions.

These and other objects, advantages, and features of the invention will become apparent to those persons  
25 skilled in the art upon reading the details of the chimeric gene, assay method, and transgenic mouse as more fully described below.

#### Brief Description of the Drawings

Figure 1 is a schematic drawing showing the  
30 construction of a chimeric MHu2M gene and a transgenic mouse containing same;

Figure 2 is a schematic view of a portion of PrP proteins showing the differences between a normal, wild-

- 15 -

type human PrP protein and a normal, wild-type mouse PrP protein;

Figure 3 shows the amino acid sequence of mouse PrP along with specific differences between mouse PrP and  
5 human PrP;

Figure 4 shows the amino acid sequence of mouse PrP and specifically shows differences between mouse PrP and bovine PrP; and

Figure 5 shows the amino acid sequence of mouse  
10 PrP and specifically shows differences between mouse PrP and ovine PrP.

#### Detailed Description of Preferred Embodiments

Before the present artificial gene, assay methodology, standardized prion preparations, and  
15 transgenic and hybrid animals used in the assay are described, it is to be understood that this invention is not limited to particular assay methods, chimeric and artificial genes, prion preparation or transgenic and hybrid animals described, as such methods, genes,  
20 preparations, and animals may, of course, vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to be limiting, since the scope of the present invention will be limited only by the  
25 appended claims.

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any methods  
30 and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, the preferred methods and materials are now described. All publications mentioned herein are incorporated herein by reference to disclose and describe



- 16 -

the methods and/or materials in connection with which the publications are cited.

### Definitions

The term "FVB" refers to a mouse strain commonly  
5 used in the production of transgenic mice. For purposes of this invention it should be noted that the mouse prion protein (PrP) gene is intact and mouse PrP is therefore expressed at normal levels.

The term "Prnp-% or Prnp-Abl" refers to a  
10 transgenic animal which has its PrP gene ablated with the "%" indicating that both alleles are ablated whereas o/+ indicates only one is ablated. Specifically, the animal being referred to is generally a transgenic mouse which has its PrP gene ablated i.e., a PrP knockout  
15 mouse. In that the PrP gene is disrupted no mouse PrP protein is expressed.

The term "sporadic CJD" abbreviated as "sCJD" refers to the most common manifestation of Creutzfeldt-Jakob Disease (CJD). This disease occurs spontaneously  
20 in individuals with a mean age of approximately 60 at a rate of 1 per million individuals across the earth.

The term "Iatrogenic CJD" abbreviated as "iCJD" refers to disease resulting from accidental infection of people with human prions. The most noted example of such  
25 is the accidental infection of children with human prions from contaminated preparations of human growth hormone.

The term "Familial CJD" refers to a form of CJD which occurs rarely in families and is inevitably caused by mutations of the human prion protein gene. The  
30 disease results from an autosomal dominant disorder. Family members who inherit the mutations succumb to CJD.

The term "Gerstmann-Strassler-Scheinker Disease" abbreviated as "GSS" refers to a form of inherited human prion disease. The disease occurs from an autosomal

- 17 -

dominant disorder. Family members who inherit the mutant gene succumb to GSS.

The term "prion" shall mean an infectious particle known to cause diseases (spongiform encephalopathies) in humans and animals. The term "prion" is a contraction of the words "protein" and "infection" and the particles are comprised largely if not exclusively of PrP<sup>Sc</sup> molecules encoded by a PrP gene. Prions are distinct from bacteria, viruses and viroids. Known prions include those which infect animals to cause scrapie, a transmissible, degenerative disease of the nervous system of sheep and goats as well as bovine spongiform encephalopathies (BSE) or mad cow disease and feline spongiform encephalopathies of cats. Four prion diseases known to affect humans are (1) kuru, (2) Creutzfeldt-Jakob Disease (CJD), (3) Gerstmann-Strassler-Scheinker Disease (GSS), and (4) fatal familial insomnia (FFI). As used herein prion includes all forms of prions causing all or any of these diseases or others in any animals used — and in particular in humans and in domesticated farm animals.

The terms "PrP gene" and "prion protein gene" are used interchangeably herein to describe genetic material which expresses proteins as shown in Figures 2-5 and polymorphisms and mutations such as those listed herein under the subheading "Pathogenic Mutations and Polymorphisms." The PrP gene can be from any animal including the "host" and "test" animals described herein and any and all polymorphisms and mutations thereof, it being recognized that the terms include other such PrP genes that are yet to be discovered.

The terms "standardized prion preparation", "prion preparation", "preparation" and the like are used interchangeably herein to describe a composition containing prions which composition is obtained from

- 18 -

brain tissue of mammals which contain substantially the same genetic material as relates to PrP proteins, e.g., brain tissue from a set of mammals which exhibit signs of prion disease which mammals may comprise any of (1) a PrP  
5 chimeric transgene; (2) have an ablated endogenous PrP gene; (3) have a high copy number of PrP genes from a genetically diverse species; or (4) are hybrids with an ablated endogenous PrP gene and a PrP gene from a genetically diverse species. The mammals from which  
10 standardized prion preparations are obtained exhibit clinical signs of CNS dysfunction as a result of inoculation with prions and/or due to developing the disease due to their genetically modified make up, e.g., high copy number of PrP genes.

15 The term "PrP gene" refers generally to any gene of any species which encodes any form of a PrP amino acid sequences including any prion protein. Some commonly known PrP sequences are described in Gabriel et al., Proc. Natl. Acad. Sci. USA 89:9097-9101 (1992) which is  
20 incorporated herein by reference to disclose and describe such sequences.

The term "artificial PrP gene" is used herein to encompass the term "chimeric PrP gene" as well as other recombinantly constructed genes which when included in  
25 the genome of a host animal (e.g., a mouse) will render the mammal susceptible to infection from prions which naturally only infect a genetically diverse test mammal, e.g., human, bovine or ovine. In general, an artificial gene will include the codon sequence of the PrP gene of  
30 the mammal being genetically altered with one or more (but not all, and generally less than 40) codons of the natural sequence being replaced with a different codon - preferably a corresponding codon of a genetically diverse mammal (such as a human). The genetically altered mammal  
35 being used to assay samples for prions which only infect

- 19 -

the genetically diverse mammal. Examples of artificial genes are mouse PrP genes encoding the sequence as shown in Figures 3, 4 and 5 with one or more different replacement codons selected from the codons shown in these Figures for humans, cows and sheep replacing mouse codons at the same position, with the proviso that not all the mouse codons are replaced with differing human, cow or sheep codons. Artificial PrP genes of the invention can include not only codons of genetically diverse animals but may include codons and codon sequences associated with genetic prion diseases such as CJD and codons and sequences not associated with any native PrP gene but which, when inserted into an animal render the animal susceptible to infection with prions which would normally only infect a genetically diverse animal.

The terms "chimeric gene," "chimeric PrP gene", "chimeric prion protein gene" and the like are used interchangeably herein to mean an artificially constructed gene containing the codons of a host animal such as a mouse with one or more of the codons being replaced with corresponding codons from a genetically diverse test animal such as a human, cow or sheep. In one specific example the chimeric gene is comprised of the starting and terminating sequence (i.e., N- and C-terminal codons) of a PrP gene of a mammal of a host species (e.g. a mouse) and also containing a nucleotide sequence of a corresponding portion of a PrP gene of a test mammal of a second species (e.g. a human). A chimeric gene will, when inserted into the genome of a mammal of the host species, render the mammal susceptible to infection with prions which normally infect only mammals of the second species. The preferred chimeric gene disclosed herein is MHu2M which contains the starting and terminating sequence of a mouse PrP gene and

- 20 -

a non-terminal sequence region which is replaced with a corresponding human sequence which differs from a mouse PrP gene in a manner such that the protein expressed thereby differs at nine residues.

5           The term "genetic material related to prions" is intended to cover any genetic material which effects the ability of an animal to become infected with prions. Thus, the term encompasses any "PrP gene", "artificial PrP gene", "chimeric PrP gene" or "ablated PrP gene"

10 which terms are defined herein as well as mutations and modifications of such which effect the ability of an animal to become infected with prions. Standardized prion preparations of the invention are produced using animals which all have substantially the same genetic

15 material related to prion so that all of the animals will become infected with the same type of prions and will exhibit signs of infection at about the same time.

          The terms "host animal" and "host mammal" are used to describe animals which will have their genome

20 genetically and artificially manipulated so as to include genetic material which is not naturally present within the animal. For example, host animals include mice, hamsters and rats which have their endogenous PrP gene altered by the insertion of an artificial gene of the

25 present invention or by the insertion of a native PrP gene of a genetically diverse test animal.

          The terms "test animal" and "test mammal" are used to describe the animal which is genetically diverse from the host animal in terms of differences between the PrP

30 gene of the host animal and the PrP gene of the test animal. The test animal may be any animal for which one wishes to run an assay test to determine whether a given sample contains prions with which the test animal would generally be susceptible to infection. For example, the

35 test animal may be a human, cow, sheep, pig, horse, cat,



- 21 -

dog or chicken, and one may wish to determine whether a particular sample includes prions which would normally only infect the test animal. This is done by including PrP gene sequences of the test animal into the host  
5 animal and inoculating the host animal with prions which would normally only infect the test animal.

The terms "genetically diverse animal" and "genetically diverse mammal" are used to describe an animal which includes a native PrP codon sequence of the  
10 host animal which differs from the genetically diverse test animal by 17 or more codons, preferably 20 or more codons, and most preferably 28-40 codons. Thus, a mouse PrP gene is genetically diverse with respect to the PrP gene of a human, cow or sheep, but is not genetically  
15 diverse with respect to the PrP gene of a hamster.

The terms "ablated prion protein gene", "disrupted PrP gene", "ablated PrP gene" and the like are used interchangeably herein to mean an endogenous prion protein gene which has been altered (e.g., add and/or  
20 remove nucleotides) in a manner so as to render the gene inoperative. Examples of non-functional PrP genes and methods of making such are disclosed in Büeler, H., et al "Normal development of mice lacking the neuronal cell-surface PrP protein" Nature 356, 577-582 (1992) which is  
25 incorporated herein by reference. Both alleles of the genes are disrupted.

The terms "hybrid animal", "transgenic hybrid animal" and the like are used interchangeably herein to mean an animal obtained from the cross-breeding of a  
30 first animal having an ablated endogenous PrP gene with a second animal which includes either (1) a chimeric gene or artificial PrP gene or (2) a PrP gene from a genetically diverse animal. For example a hybrid mouse is obtained by cross-breeding a mouse with an ablated  
35 mouse PrP gene with a mouse containing (1) human PrP



- 22 -

genes (which may be present in high copy numbers) or (2) chimeric genes. The term hybrid includes any offspring of a hybrid including inbred offspring of two hybrids provided the resulting offspring is susceptible to  
5 infection with prions with normal infect only a genetically diverse species.

The terms "susceptible to infection" and "susceptible to infection by prions" and the like are used interchangeably herein to describe a transgenic or  
10 hybrid test animal of the invention which develops a prion disease if inoculated with prions which would normally only infect a genetically diverse test animal. The terms are used to describe a transgenic or hybrid animal of the invention such as a transgenic mouse  
15 Tg(MHu2M) which, without the chimeric PrP gene, would not be susceptible to infection with a human prion (less than 20% chance of infection) but with the chimeric gene is susceptible to infection with human prions (80% to 100% chance of infection).

20 The term "incubation time" shall mean the time from inoculation of an animal with a prion until the time when the animal first develops detectable symptoms of disease resulting from the infection. A reduced incubation time is one year or less, preferable about  
25 200 days  $\pm$  50 days or less, more preferably about 50 days  $\pm$  20 days or less.

Abbreviations used herein include:

CNS for central nervous system;  
BSE for bovine spongiform encephalopathy;  
30 CJD for Creutzfeldt-Jakob Disease;  
FFI for fatal familial insomnia;  
GSS for Gerstmann-Strassler-Scheinker Disease;  
Hu for human;  
HuPrP for a human prion protein;  
35 Mo for mouse;

- 23 -

- MoPrP for a mouse prion protein;  
SHa for a Syrian hamster;  
SHaPrP for a Syrian hamster prion protein;  
Tg for transgenic;
- 5 Tg(SHaPrP) for a transgenic mouse containing the PrP gene  
of a Syrian hamster;  
Tg(HuPrP) for transgenic mice containing the complete  
human PrP gene;  
Tg(ShePrP) for transgenic mice containing the complete  
10 sheep PrP gene;  
Tg(BovPrP) for transgenic mice containing the complete  
cow PrP gene;  
PrP<sup>Sc</sup> for the scrapie isoform of the prion protein;  
MoPrP<sup>Sc</sup> for the scrapie isoform of the mouse prion  
15 protein;  
MHu2M for a chimeric mouse/human PrP gene wherein a  
region of the mouse PrP gene is replaced by a  
corresponding human sequence which differs from  
mouse PrP at 9 codons;
- 20 Tg(MHu2M) mice are transgenic mice of the invention which  
include the chimeric MHu2M gene;  
MHu2MPrP<sup>Sc</sup> for the scrapie isoform of the chimeric human/  
mouse PrP gene;  
PrP<sup>CJD</sup> for the CJD isoform of a PrP gene;
- 25 Prn-p<sup>0/0</sup> for ablation of both alleles of an endogenous  
prion protein gene, e.g., the MoPrP gene;  
Tg(SHaPrP<sup>+/0</sup>)81/Prn-p<sup>0/0</sup> for a particular line (81) of  
transgenic mice expressing SHaPrP, +/0 indicates  
heterozygous;
- 30 Tg(HuPrP)/Prnp<sup>0/0</sup> for a hybrid mouse obtained  
by crossing a mouse with a human prion protein gene  
(HuPrP) with a mouse with both alleles of the  
endogenous prion protein gene disrupted;  
Tg(MHu2M)/Prnp<sup>0/0</sup> for a hybrid mouse obtained

- 24 -

by crossing a mouse with a chimeric prion protein gene (MHu2M) with a mouse with both alleles of the endogenous prion protein gene disrupted.

FVB for a standard inbred strain of mice often used  
5 in the production of transgenic mice since eggs of FVB mice are relatively large and tolerate microinjection of exogenous DNA relatively well.

#### General Aspects of the Invention

The present invention includes several aspects  
10 including: (1) an artificial gene comprised of codon sequences which when inserted into the genome of a host animal (e.g. a mouse or hamster) will render the animal susceptible to infection with prions which normally infect only a genetically diverse test animal (e.g. a  
15 human, cow or sheep), thereby including genes wherein one or more codons of a naturally occurring PrP gene of a host animal are replaced with corresponding codons of a genetically diverse test animal; (2) a chimeric gene which gene is comprised of the codon sequence of a PrP  
20 gene of a host mammal of a first species (preferably a mouse or hamster) which gene has been modified to include a corresponding segment of a PrP gene of a test animal, preferably a mammal such as a human, cow or sheep; (3) a transgenic mammal containing an artificial gene of the  
25 invention such as a transgenic mouse including a chimeric PrP gene wherein a portion of the mouse gene is replaced with a corresponding portion of a human PrP gene thereby rendering the mouse susceptible to infection with human prions; (4) a transgenic mammal with elevated levels of  
30 expression of a PrP gene of a genetically diverse mammal wherein the elevated levels of expression are obtained by incorporating a high copy number (30 or more) of a native PrP gene of a genetically diverse test animal and/or the inclusion of an enhanced promoter operatively fused to

- 25 -

the PrP gene of a genetically diverse animal; (5) a transgenic hybrid animal which is obtained by crossing a animal having an ablated endogenous PrP gene with an animal with a chimeric gene as per (2) above or an animal  
5 with a PrP gene of another genetically diverse animal therein e.g., as per (4) above; (6) standardized prion preparations which contain the same amount (preferably at the same concentration) and type of prions in each preparation; (7) a method of determining whether a sample  
10 is infected with prions which method involves inoculating a transgenic or hybrid mammal of the invention with a sample to be tested (and preferably simultaneously inoculating identical test animals with a standardized prion preparation for use as controls) and observing the  
15 mammal(s) for a period of time sufficient to determine if the mammal(s) develop(s) symptoms of a disease normally associated with prions; (8) a method of testing the efficacy of a drug in the treatment of disease developed as a result of infection with prions comprising  
20 administering a drug to be tested to a transgenic or hybrid animal infected with prions (preferably a standardized prion preparation) and observing and/or testing the mammal to determine if the drug aids in treating or slowing the progress of the disease or its  
25 symptoms; and (9) a method for determining the cause of death of an animal comprising inoculating a transgenic or hybrid animal of the invention with body fluid or tissue such as extracted brain tissue from the animal which has died (and preferably inoculating control animals with a  
30 standardized preparation of prions) and observing the transgenic or hybrid animal (and control animals) in order to determine if the animal(s) develop(s) symptoms of prion infections.

- 26 -

Preferred host animals are mice and hamsters, with mice being most preferred in that there exists considerable knowledge on the production of transgenic animals. Other possible host animals include those  
5 belonging to a genus selected from *Mus* (e.g. mice), *Rattus* (e.g. rats), *Oryctolagus* (e.g. rabbits), and *Mesocricetus* (e.g. hamsters) and *Cavia* (e.g., guinea pigs). In general mammals with a normal full grown adult body weight of less than 1 kg which are easy to breed and  
10 maintain can be used. The host PrP gene can be changed to include codons from genetically diverse PrP genes from test animals belonging to a genus selected from *Bos*, *Ovis*, *Sus* and *Homo*. Preferably, a mouse host PrP gene is changed to include codons from a human, cow or sheep PrP  
15 gene, with human being most preferred. Humans are preferred because an important object of the invention is to use the animal to test a sample of material to determine if that material has prions which will infect a human and cause a human to develop a CNS disease such as  
20 CJD.

The genetic material which makes up the PrP gene is known for a number of different species of animals (see Gabriel et al., Proc. Natl. Acad. Sci. USA 89:9097-9101 (1992)). Further, there is considerable homology between  
25 the PrP genes in different mammals. For example, see the amino acid sequence of mouse PrP compared to human, cow and sheep PrP in Figures 3, 4 and 5 wherein only the differences are shown. Further, note that the segment of a PrP gene used to create the MHu2M gene of the present  
30 invention will result in encoding of protein which shows a difference between the human and a mouse protein of only nine residues. Although there is considerable genetic homology with respect to PrP genes, the differences are significant in some instances. More  
35 specifically, due to small differences in the protein

- 27 -

encoded by the PrP gene of different mammals, a prion which will infect one mammal (e.g. a human) will not normally infect a different mammal (e.g. a mouse). Due to this "species barrier", it is not generally possible to use normal animals, (i.e., animal which have not had their genetic material related to prions manipulated) such as mice to determine whether a particular sample contains prions which would normally infect a different species of animal such as a human. The present invention solves this problem in a surprising manner.

Relationships - PrP genes: copy numbers: genetic diversity

Commercially useful transgenic animals are preferably small and easy to reproduce; thus, host animals such as mice, hamsters, guinea pigs and rats are preferred, with mice being most preferred. In order for the transgenic animals to be useful, it is necessary for the animals to be susceptible to infection with prions which normally infect only genetically diverse test animals, and in particular animals of commercial significance for testing, such as humans, cows, horses, sheep, pigs, cats, dogs and chickens, with humans being most preferred. Further, for the transgenic and hybrid animals to be useful in a practical and commercial sense, it is necessary for the animals to demonstrate symptoms of the disease within a relatively short period after inoculation, and for a very high percentage of the animals to demonstrate symptoms of the disease after inoculation e.g., 80% or more of inoculated animals develop clinical symptoms of CNS disease with 350 days or less after inoculation.

In producing a transgenic animal having the above-described characteristics, we noted a number of relationships of significance. First, when the entire PrP gene of a test animal (such as a human) is made



- 28 -

functional in the host animal (such as a mouse) the resulting transgenic animal (with a low copy number of human PrP genes) is not susceptible to infection with human prions.

5       Second, we found that infection would occur if the endogenous PrP gene of the host animal is ablated.

Third, when only some of the codons differing between the host and the test animal are switched, the resulting transgenic animal is susceptible to infection  
10 with prions which normally only infect the test animal.

Fourth, we noticed that, as the copy number of the artificial gene in the transgenic animal is increased, the incubation time in some cases decreases.

Fifth, we noted that humans with some genetic  
15 defects resulting in prion diseases have different genetic defects in their PrP gene and that by matching the defects in any transgenic animal will render that animal more susceptible to infection with prions from the diseased human.

20       With this knowledge, we deduced that it is possible to produce a transgenic animal wherein all of the PrP gene of the host animal is replaced with the PrP gene of a test animal to obtain a useful transgenic animal which is susceptible to infection with prions which normally  
25 only infect the test animal by substantially increasing the copy number of the test animal's PrP gene in the host animal and preferably also ablating the endogenous PrP gene. For example, a transgenic mouse which includes the entire PrP gene of a human in a relatively low copy  
30 number (e.g. 1 to 4) is not susceptible to infection with human prions (unless the endogenous mouse PrP gene is ablated). However, if the transgenic mouse includes a very high copy number of a human gene (e.g. 30-300 copies), the resulting transgenic animal is susceptible  
35 to infection with human prions. Further, when a host

- 29 -

animal such as a mouse has only a portion of its PrP gene replaced with a corresponding portion of a test animal such as a human, the resulting transgenic animal is highly susceptible to infection with prions which

5 normally infect only the test animal. This is true even if the chimeric gene is present in the transgenic animal in a relatively low copy number (e.g. 1 to 4 copies) resulting in low expression of MHu2M PrP<sup>C</sup>.

Lastly, in order to reduce incubation time hybrid  
10 mice were created by crossing mice with ablated PrP genes with transgenic mice which (1) included a PrP gene from a genetically diverse animal e.g., a human or (2) include a chimeric or artificial gene of the present invention. The chimeric gene and/or PrP gene from genetically  
15 diverse test animal may be present in high copy number. The copy number can be increased in order to reduce incubation time provided the copy number is not increased so far that the animal becomes spontaneously ill, i.e., become ill without inoculation with prions.

20 Based on the above, it can be understood that the preferred transgenic animals of the invention are (1) animals such as mice which include a chimeric PrP gene, i.e., only a portion, but not all, of their PrP gene replaced with a corresponding portion of the PrP gene of  
25 a test animal or (2) animals with an ablated endogenous PrP gene and a PrP gene from another animal such as a human most preferable where that human PrP gene has a genetic defect which results in a prion disease when in a human.

30 It is preferable to include chimeric genes within the transgenic animal in a relatively high copy number, in that increasing the copy number tends to decrease the incubation time for the disease once the animal is inoculated with material containing prions.

35 Notwithstanding such, we now understand that, when the

- 30 -

copy number is increased to very high numbers (e.g. 100 copies and above), the transgenic animals may spontaneously demonstrate symptoms of prion disease. Thus, a most preferred transgenic animal of the invention will include a chimeric PrP gene in a sufficiently high copy number so as to shorten the incubation time (e.g. 50 copies  $\pm$  25) but in a sufficiently low number so as to not initiate spontaneous symptoms characteristic of prion diseases (e.g., not more than 100 copies). It will be understood by those skilled in the art that the number of copies necessary in order to obtain elevated levels of expression of the PrP gene will vary depending upon the particular gene inserted into the particular host. Adjustments can be made to reduce the copy number if the resulting transgenic animals become spontaneously ill. Alternatively adjustments can be made to increase the copy number if the resulting transgenic animals are not subject to infection with prions which normally infect only a genetically diverse animal. Further, adjustments can be made with respect to the use of specific types of enhanced promoters in order to elevate the levels of expression without increasing copy numbers. Specific types of enhanced promoters are known such as neuronal enolase promoters which would provide enhanced expression to the PrP gene without increased copy numbers. The enhanced promoters may operate constitutively or inducibly.

The ability to successfully produce a transgenic animal is related, in part, to the genetic diversity between the host animal and the test animal as regards their respective PrP genes. For example, the PrP gene of a mouse and a hamster are relatively similar in that they differ only at 16 codons out of a total of 254 codons. When the genetic similarity of the PrP genes are this close, it is possible to include the entire PrP gene

- 31 -

sequence of the test animal into the host animal and render the host animal susceptible to prions which normally only infect the test animal even with the host animals endogenous PrP intact i.e., not ablated.

- 5 However, such is not the case when the host animal and test animal are genetically diverse, i.e. differ in PrP genes by 20 or more codons. Thus, when a mouse PrP gene is completely replaced with a genetically diverse PrP gene, such as that of a human, the resulting transgenic
- 10 mouse will not be susceptible to infection with human prions unless (1) the endogenous PrP gene of the mouse is ablated or (2) the human gene is present in the mouse in a relatively high copy number -- which high copy number may result in spontaneous development of disease and/or
- 15 (3) the human PrP gene includes a genetic defect which results in a prion disease in a human.

To solve the problem of being able to decrease the copy number such that the animal would not spontaneously become sick, and yet allow the animal to become sick when

20 inoculated with human prions, we created a chimeric gene which includes only a portion of the human PrP gene in the mouse PrP gene. A more specific description of how the species barrier was broken in accordance with the present invention is provided below.

- 25 When transgenic animals (with endogenous PrP gene intact) are produced by placing the entire human prion protein gene into that of a mouse the resulting transgenic mouse does not become consistently ill in a short period of time when inoculated with prions which
- 30 generally only infect humans i.e., is not susceptible to infection with human prions. The inability to become infected appears to be related to the presence of the endogenous mouse prion protein gene. When a mouse with a human prion protein gene is crossed with a mouse with a
- 35 disrupted endogenous mouse PrP gene the hybrid offspring

- 32 -

are infected by prions which normally only infect humans. Such hybrid mice will consistently become infected and exhibit an incubation time of less than 300 days, preferably 250 or less  $\pm$  50 days.

#### 5 Species Barrier Broken

The transmission of human CJD to apes and monkeys 1.5 - 3 years after intracerebral inoculation provided considerable interest in the causes of neurodegenerative diseases [Gibbs, Jr. et al., Science 161:388-389 (1968)].

10 Humans are not genetically diverse from apes and monkeys which accounts for the cross-species infectivity, although with a long incubation time. While the high cost of caring for nonhuman primates prevented extensive studies of the human prion diseases, the transmissibility  
15 of these diseases stimulated studies of the animal prion analogues in rodents [Manuelidis et al., Proc. Natl. Acad. Sci. USA 75:3422-3436 (1978); Manuelidis et al., Proc. Natl. Acad. Sci. USA 73:223-227 (1976); Tateishi et al., Ann. Neurol. 5:581-584 (1979)].

20 The present disclosure opens a new frontier in the investigation of the human prion diseases since transmission studies can now be performed relatively rapidly in genetically altered mammals such as Tg(MHu2M) mice that are relatively inexpensive to maintain. For  
25 the first time, endpoint titrations of prions in multiple human body tissues and fluids can be performed and standard curves constructed for more economical incubation time assays. The information derived from such studies of human prions will be useful in the  
30 management of CJD patients who are thought to pose some risk to relatives, physicians, nurses and clinical laboratory technicians [Berger et al., Neurology 43:205-206 (1993); Ridley et al., Lancet 341:641-642 (1993)].

- 33 -

In studies of human prion diseases with apes and monkeys, the use of one or two, or rarely three, animals as recipients for a single inoculum has presented a significant problem in evaluating the transmissibility of a particular inoculum from an individual patient. The transgenic mice contain a chimeric prion protein gene, e.g., Tg(MHu2M) mice, and hybrid mice e.g., Tg(HuPrP)/Prnp<sup>0/0</sup> described here obviate many of the problems created by using nonhuman primates.

10        These results demonstrate the "universality" of the MHu2M transgene for transmission studies with other types of transgenic animals and other prion inocula. For example, it may be most efficient to use mice expressing MHu2MP<sub>PrP</sub> transgenes coding for either a methionine or  
15 valine at codon 129, and by doing so, match the genotype of the Tg mouse (with respect to codon 129) with the genotype of the individual from which the inoculum is derived. Homozygosity at the codon 129 polymorphism has a profound influence on the incidence of sporadic CJD  
20 [Palmer et al., Nature 352:340-342 (1991)]. The MHu2MP<sub>PrP</sub> transgene encodes a Met at codon 129 and the iatrogenic CJD case was homozygous for Met [Collinge et al., Lancet 337:1441-1442 (1991)].

A human PrP gene is polymorphic at codon 129. More specifically, normal human PrP gene can be either  
25 homozygous Met/Met or Val/Val or heterozygous Met/Val at codon 129. The codon 129 polymorphism influences the susceptibility of humans to prion disease and specifically to iatrogenic and sporadic CJD. This  
30 polymorphic codon is contained in the central region of MHu2MP<sub>PrP</sub> which is derived from human PrP. The DNA sequence used to generate Tg(MHu2M) mice encodes Met at codon 129. The transgenic mice expressing MHu2MP<sub>PrP</sub> with valine at codon 129 can be produced using similar  
35 procedures.



- 34 -

To break the species barrier we have created an artificial PrP gene which, when inserted into a host mammal (such as a mouse) renders that mammal susceptible to infection with prions which normally infect only a genetically diverse test mammal (e.g. a human, cow or sheep). The artificial PrP gene may include the natural PrP gene sequence of the host animal with one or more (preferably less than 40) codon sequences being replaced with other codon sequences such as corresponding codons of a genetically diverse mammal (e.g. a human, cow or sheep).

In a specific example of the invention the species barrier is broken by inserting into a mammal (a mouse) the chimeric gene (MHu2M) which is shown being assembled schematically in Figure 1. In order to produce the chimeric gene, it is first necessary to obtain nucleotide sequences which encode human PrP. The human PrP genes are then subjected to the conventional PCR procedures in order to produce large numbers of copies of the gene or portions of the gene. The PCR product is then isolated, specific restriction sites are added and the copied product is subjected to specific endonucleases in order to remove a middle section of the human PrP gene. Specifically, restriction sites are added such that when the PCR product is subjected to endonucleases such as Asp718 as well as BstEII, a section of the gene is cut out. The use of these two endonucleases will remove a center portion of the human PrP gene (codons 94-188) which portion encodes amino acid residues 94 through 188. Endonucleases are also used to remove a corresponding center portion of the mouse PrP gene. The removed center portion of the mouse gene is then discarded and the center portion obtained from the human PrP gene is fused into the mouse gene to produce a chimeric human/mouse gene. Details of how the specific MHu2M gene was

- 35 -

produced are described in Example 1 and shown in Figure 1.

As shown with Figure 2, there is a high degree of homology between the removed center portion of the human PrP gene and the segment of the mouse PrP gene which is replaced. Specifically, the segments differ at nine codons. Thus, when the genetic material is expressed, the resulting chimeric MHu2M protein will differ from MoPrP at 9 residues. These residues and their positions are shown in Figure 2. After the chimeric gene is produced, it can be microinjected into a mouse egg using known technology as described within Scott et al., Cell 59:847-857 (1989) and Scott et al., Protein Sci. 1:986-997 (1992) and see also WO91/19810 published December 22, 1991 as well as other publications relating to the production of transgenic mice cited therein and known to those skilled in the art. The injected mouse egg is then implanted into a mouse using known procedures. Multiple eggs can be implanted into a single mouse and known procedures can be used to determine whether the resulting offspring are transgenic mice which include the chimeric gene within their genome. Details of this procedure are described in Example 1.

We have successfully broken the "species barrier" by producing a chimeric PrP gene wherein a middle portion of the mouse PrP gene is replaced with a corresponding middle portion of a human PrP gene thereby leaving the C- and N-terminus of the mouse PrP gene intact. However, other segments of the mouse PrP gene can be replaced with other homologous segments of the human PrP gene and obtain a transgenic mouse which is subject to being readily infected with human prions. Thus, the invention is not limited to the particular chimeric gene MHu2M or chimeric mice produced using this gene. The invention includes all types of transgenic animals which include

- 36 -

artificial genes wherein the artificial gene renders the transgenic animal susceptible to infection with prions which normally infect only a genetically diverse animal.

#### Additional Chimeric Constructs

5       The chimeric construct MHu2M described above includes codons for human PrP gene in the middle section of a mouse PrP gene. In that positive results were obtained using such a chimeric gene three additional chimeric genes were produced which are referred to a  
10 MHu3, Hu3M and Hu4M. The chimeric gene MHu3 is comprised of a mouse PrP sequence from the start codon of the PrP open reading frame up until codon 94 and has a human PrP sequence from codon 94 to the end of the PrP open reading frame. The chimeric construct Hu3M is comprised of a  
15 human PrP sequence from the start of the PrP open reading frame until codon 188 followed by a mouse PrP sequence from codon 188 to the end of the PrP open reading frame. The chimeric construct Hu4M is comprised of a human PrP sequence from the start of the PrP open reading frame up  
20 until codon 223 followed by a mouse PrP sequence from codon 223 to the end of the PrP open reading frame.

Numerous specific examples of artificial genes of the invention can be deduced from reviewing Figures 3, 4 and 5. Specifically, one may start with the basic PrP  
25 gene of a mouse (as the host animal) which animal is to become the transgenic animal. Thereafter, one or more codons of the mouse gene may be replaced with one or more corresponding codons of a human, bovine or sheep PrP gene which codon is different from the corresponding codon of  
30 the mouse gene but at the same relative position in the gene. By showing that it is possible to break the "species barrier" by creating a particular chimeric gene whereby transgenic mice can test for the presence of human prions we have opened the door for the creation of

- 37 -

other transgenic animals which will include other artificial PrP genes which, for example, can allow for the testing for the presence of bovine or ovine prions in a sample. The chimeric or artificial PrP genes can be  
5 used by themselves or in an animal with an ablated PrP gene background.

#### High Copy Numbers

The present invention includes transgenic animals wherein the host animal has its genome changed to include  
10 multiple copies of the entire PrP gene of a genetically diverse test animal. Thus, for example, the invention includes transgenic mice and hamsters altered to include two fold or higher levels of expression of the PrP gene of a genetically diverse test animal such as a human, cow  
15 or sheep. The two fold or higher levels of expression can be obtained by including higher copy numbers such as 30 or more copies of the PrP gene of the genetically diverse test animal and/or by including an enhanced promoter which elevates the level of expression of even a  
20 low copy number of the gene.

#### Hybrid Animals

Hybrid animals of the invention can be produced by crossing an animal with an ablated endogenous prion protein gene with either of the transgenic animals  
25 mentioned above. For example, a mouse containing a human/mouse chimeric prion is crossed with a mouse with a disrupted endogenous prion protein gene e.g., Tg(Prnp<sup>0/0</sup>). Alternatively, a mouse containing a high copy number of human prion protein genes (e.g., 50 ± 25) is crossed with  
30 a mouse with a disrupted endogenous prion protein gene e.g., Tg(Prnp<sup>0/0</sup>) to obtain a hybrid Tg(HuPrP)/Prnp<sup>0/0</sup>. A variety of different hybrids can be obtained by crossing an animal with an ablated prion protein gene (i.e., a

- 38 -

null prion background) with different transgenic animals with different prion protein genes. When successful hybrids are obtained they can be crossed to produce other animals which for the purpose of the disclosure are also  
5 considered hybrids if they are susceptible to infection with prions which generally only infect a genetically diverse species. A null prion background means that more than 50% of the endogenous prion protein genes are disrupted, preferable more than 80%, more preferable more  
10 than 90% and most preferable 100% so that no endogenous PrP is expressed.

Tg(MHu2M) mice with shorter incubation times

The incubation time of Tg(MHu2M) mice inoculated with Hu prions is now about 200 days or less  $\pm 50$  days,  
15 which can be reduced substantially by increasing the copy number of the MHu2M gene (e.g. to about  $50 \pm 25$ ) and thereby obtaining an elevated level of expression. In Tg(SHaPrP) mice, the level of SHaPrP transgene expression was found to be inversely proportional to the length of  
20 the scrapie incubation time after inoculation with SHA prions [Prusiner et al., Cell 63:673-686 (1990)]. Thus, producing Tg(MHu2M) mice with higher levels of transgene expression is a means of substantially reducing incubation time.

25 Based on other studies with Tg mice harboring a hamster PrP gene we have shown that the incubation period for Tg mice (or any host) with a chimeric or artificial PrP gene can approach 30 days or less. For example, Tg(SHaPrP)/Prnp<sup>0/0</sup> mice containing more than 100 copies of  
30 a exogenous PrP gene material have an incubation time of approximately 25 days.

In addition, removing MoPrP<sup>c</sup> by crossing Tg(MHu2M) mice onto an ablated background (Prn-p<sup>0/0</sup>) may also reduce the incubation time since Tg(SHaPrP<sup>0/0</sup>)81/Prn-p<sup>0/0</sup> mice

- 39 -

exhibit a 30% reduction in incubation times compared to Tg(SH<sub>2</sub>PrP<sup>0</sup>)<sup>81</sup>/Prn-p<sup>+/+</sup> mice [Büeler et al., Cell 73:1339-1347 (1993). Prusiner et. al., Proc. Natl. Acad. Sci. USA 90:10608-10612 Nov. 1993. Accordingly, we have also  
5 used fertilized eggs from mice in which the endogenous PrP gene has been ablated as recipients for microinjection of the MHu2M PrP construct.

By systematically altering the extent and position of the substitutions in other chimeric Hu/Mo PrP  
10 constructs, it is possible to further enhance the susceptibility of Tg mice to Hu prions as reflected by shortened incubation times. Shortening the incubation time is a worthwhile goal for the facilitation of many future studies in prion research and for the evaluation  
15 of pharmaceuticals, foods, tissues, organs, grafts, cosmetics and other substances - particularly substances which have some portion derived from an animal, such as a human, which animal might be infected with prions.

In general, there is an inverse relationship between  
20 the number of copies of a chimeric or artificial PrP gene and the incubation time of disease after inoculation of the transgenic animal with prions. Specific MHu2M mice disclosed herein have only 3 or 4 copies of the MHu2M gene. The number of copies can be increased to 30 to  
25 400, thereby significantly reducing the incubation time from about 200 days to 50 days ± 20 days or less. Those skilled in the art will understand that the copy number should not be increased to a point where the animal will spontaneously develop disease.

### 30 Pathogenic mutations and polymorphisms

There are a number of known pathogenic mutations in the human PrP gene. Further, there are known polymorphisms in the human, sheep and bovine PrP genes.



- 40 -

The following is a list of such mutations and polymorphisms:

MUTATION TABLE

	Pathogenic human mutations	Human Polymorphism s	Sheep Polymorphism s	Bovine Polymorphisms
5	2 octarepeat insert	Codon 129 Met/Val	Codon 171 Arg/Glu	5 or 6 octarepeats
10	4 octarepeat insert	Codon 219 Glu/Lys	Codon 136 Ala/Val	
	5 octarepeat insert			
	6 octarepeat insert			
15	7 octarepeat insert			
	8 octarepeat insert			
20	9 octarepeat insert			
	Codon 102 Pro-Leu			
	Codon 105 Pro-Leu			
25	Codon 117 Ala-Val			
	Codon 145 Stop			
	Codon 178			
30	Asp-Asn			
	Codon 180 Val-Ile			
	Codon 198 Phe-Ser			
35	Codon 200 Glu-Lys			
	Codon 210 Val-Ile			
	Codon 217			
40	Asn-Arg			
	Codon 232 Met-Ala			

- 41 -

Gene Constructs Containing Mutations

Several different PrP gene constructs are described above. One typical example of the constructs is MHu2M which includes the N- terminal and C- terminal portions of a mouse PrP gene but a center portion replaced with a human PrP gene wherein the human sequence includes codons corresponding to the mouse codons replaced. As indicated above similar chimeric genes are produced by replacing corresponding codons within the middle area of the mouse PrP gene with codons of the PrP gene of a sheep or cow. The mouse produced would then be susceptible, respectively, to prions which infect sheep or cows. However, this general method of creating chimeric genes and using the genes to create chimeric mice can be further refined as described below.

The above mutation table shows that many animals include specific mutations and polymorphisms at specific sites. When a mutation occurs the individual with the mutation may develop the symptoms of prion disease without ingesting infectious prions. Homogenized brain tissue from such individuals can be used to inoculate transgenic animals that will then develop symptoms of prion disease.

Transgenic mice with chimeric PrP genes inoculated with brain tissue derived from an individual with prion disease resulting from a mutation per the above mutation table may not cause symptoms or may cause symptoms of prion disease only after an extended time. The present invention provides a means of resolving this problem. Specifically, the chimeric gene is created using a sequence from another animal which includes a mutation. As an example the human middle section can include the human mutation at codon 102. When the human sequence portion is included with the 102 mutation and a transgenic animal is created the resulting transgenic

- 45 -

of sources was less than 10% in Tg(HuPrP) mice, no different from the rate observed in non-Tg mice. Likewise the conversion of HuPrP<sup>C</sup> into HuPrP<sup>Sc</sup> in Tg(HuPrP) mice appears to be a relatively infrequent event similar to the rare conversion of MoPrP<sup>C</sup> to PrP<sup>Sc</sup> in response to human prions. The low rates of transmission in these mice do not seem to be a consequence of low titers of human prion titers: two inocula which failed to cause disease in Tg(HuPrP) mice transmitted to 100% of inoculated Tg(MHu2M) animals.

#### New Approaches To Investigating Human Prion Diseases

The remarkable sensitivity of Tg(MHu2M) mice to Hu prions represents an important advance in neurodegenerative disease research. Based on the present disclosure regarding chimeric Hu/Mo PrP transgenes we conceived of a similar approach to the construction of Tg mice susceptible to BSE and scrapie sheep prions. Such would be useful in detecting prion diseases in domestic animals. The importance of animal prion diseases is illustrated by BSE or "mad cow disease" in Great Britain, where >150,000 cattle have died and serious consideration has been given to slaughtering millions of cattle potentially infected with prions. This prion disease BSE is thought to have originated with cattle consuming meat and bone meal produced from sheep offal containing scrapie prions [Wilesmith, J.W., Semin. Viro. 2:239-245].

The BSE epidemic has led to considerable concern about the safety for humans of European beef and other cattle products. Epidemiologic studies over the past two decades have provided much data arguing that humans are unlikely to contract CJD from scrapie-infected sheep products [Harries-Jones et al., J. Neurol. Neurosurg. Psychiatry 51:1113-1119 (1988); Cousens et al., J. Neurol. Neurosurg. Psychiatry 53:459-465 (1990); Brown

- 46 -

et al., Neurology 37:895-904 (1987)]. There are seven amino acid substitutions which distinguish bovine from sheep PrP which must be considered in drawing conclusions from sheep scrapie about the risk factors to humans from BSE. Whether any of these seven amino acid substitutions render bovine prions permissive in humans remains to be established. It may be that Tg(MHu2M) mice are susceptible to bovine as well as sheep prions. Of perhaps even greater importance, Tg(MHu2M) mice have immediate application in the testing of pharmaceuticals for human prion contamination. The Tg(MHu2M) mice described here provide a sensitive, reliable and economical bioassay for detecting the presence of human prions.

#### 15 Standardized Prion Preparation

Standardized prion preparations are produced for use in assays so as to improve the reliability of the assay. Although the preparation can be obtained from any animal it is preferably obtained from a host animal which has brain material containing prions of a test animal. For example, a Tg mouse containing a human prion protein gene can produce human prions and the brain of such a mouse can be used to create a standardized human prion preparation. The preparation can be further standardized by repeating the above process. More specifically, per the above process some prion containing material must be used to inoculate the transgenic mice. The source of that prion containing material may itself be unpredictable and result in infecting transgenic mice in different ways. Thus, if the transgenic mice are infected with a non-standard material some may develop the symptoms of prion disease at different rates and some may not develop symptoms at all. If a group of mice which develops symptoms of prion disease at the same time

- 47 -

are sacrificed and their brains extracted and homogenized such will create a relatively standard prion preparation. This preparation can then be used to inoculate a new group of transgenic animals. This process can be  
5 repeated a number of times e.g., 1 to 10 times or until such point as all of the transgenic mice are developing symptoms of prion disease at approximately the same point in time after inoculation with the standardized preparation. Further details of how to produce a  
10 standardized preparation are provided below.

In that the preparation is to be a "standard" it is preferably obtained from a battery (e.g., 100; 1,000, or more animals) of substantial identical animals. For example, 100 mice all containing a very high copy number  
15 of human PrP genes (all polymorphisms and mutations) would spontaneously develop disease and the brain tissue from each could be combined to make a useful standardized prion preparation.

Standardized prion preparations can be produced  
20 using any of the modified host mammals of the present invention. For example, standardized prion preparations could be produced using mice, rats, hamsters, or guinea pigs which are genetically modified per the present invention so that they are susceptible to infection with  
25 prions which prions would generally only infect genetically diverse species such as a human, cow, sheep or horse and which modified host mammals will develop clinical signs of CNS dysfunction within a period of time of 350 days or less after inoculation with prions. The  
30 most preferred host mammal is a mouse in part because they are inexpensive to use and because a greater amount of experience has been obtained with respect to production of transgenic mice than with respect to the production of other types of host animals.

- 48 -

Once an appropriate type of host is chosen, such as a mouse, the next step is to choose the appropriate type of genetic manipulation to be utilized to produce a standardized prion formulation. For example, the mice  
5 may be mice which are genetically modified by the insertion of a chimeric gene of the invention. Within this group the mice might be modified by including high copy numbers of the chimeric gene and/or by the inclusion of multiple promoters in order to increase the level of  
10 expression of the chimeric gene. Alternatively, hybrid mice of the invention could be used wherein mice which have the endogenous PrP gene ablated are crossed with mice which have a human PrP gene inserted into their genome. There are, of course, various subcategories of  
15 such hybrid mice. For example, the human PrP gene may be inserted in a high copy number and/or used with multiple promoters to enhance expression. In yet another alternative the mice could be produced by inserting multiple different PrP genes into the genome so as to  
20 create mice which are susceptible to infection with a variety of different prions, i.e., which generally infect two or more types of test animals. For example, a mouse could be created which included a chimeric gene including part of the sequence of a human, a separate chimeric gene  
25 which included part of the sequence of a cow and still another chimeric gene which included part of the sequence of a sheep. If all three different types of chimeric genes were inserted into the genome of the mouse the mouse would be susceptible to infection with prions which  
30 generally only infect a human, cow and sheep.

After choosing the appropriate mammal (e.g., a mouse) and the appropriate mode of genetic modification (e.g., inserting a chimeric PrP gene) the next step is to produce a large number of such mammals which are  
35 substantially identical in terms of genetic material



- 49 -

related to prions. More specifically, each of the mice produced will include an identical chimeric gene present in the genome in substantially the same copy number. The mice should be sufficiently identical genetically in  
5 terms of genetic material related to prions that 95% or more of the mice will develop clinical signs of CNS dysfunction within 350 days or less after inoculation and all of the mice will develop such CNS dysfunction at approximately the same time e.g., within  $\pm$  30 days of  
10 each other.

Once a large group e.g., 50 or more, more preferably 100 or more, still more preferably 500 or more of such mice are produced. The next step is to inoculate the mice with prions which generally only infect a  
15 genetically diverse mammal e.g., prions from a human, sheep, cow or horse. The amounts given to different groups of mammals could be varied. After inoculating the mammals with the prions the mammals are observed until the mammals exhibit symptoms of prion infection e.g.,  
20 clinical signs of CNS dysfunction. After exhibiting the symptoms of prion infection the brain or at least a portion of the brain tissue of each of the mammals is extracted. The extracted brain tissue is homogenized which provides the standardized prion preparation.

25 As an alternative to inoculating the group of transgenic mice with prions from a genetically diverse animal it is possible to produce mice which spontaneously develop prion related diseases. This can be done, for example, by including extremely high copy numbers of a  
30 human PrP gene into a mouse genome. When the copy number is raised to, for example, 100 or more copies, the mouse will spontaneously develop clinical signs of CNS dysfunction and have, within its brain tissue, prions which are capable of infecting humans. The brains of  
35 these animals or portions of the brain tissue of these

- 50 -

animals can be extracted and homogenized to produce a standardized prion preparation.

The standardized prion preparations of the invention can be used directly or can be diluted and tittered in a manner so as to provide for a variety of different positive controls. More specifically, various known amounts of such standardized preparation can be used to inoculate a first set of transgenic control mice. A second set of substantially identical mice are inoculated with a material to be tested i.e., a material which may contain prions. A third group of substantially identical mice are not injected with any material. The three groups are then observed. The third group, should, of course not become ill in that the mice are not injected with any material. If such mice do become ill the assay is not accurate probably due to the result of producing mice which spontaneously develop disease. If the first group, injected with a standardized preparation, do not become ill the assay is also inaccurate probably because the mice have not been correctly created so as to become ill when inoculated with prions which generally only infect a genetically diverse mammal. However, if the first group does become ill and the third group does not become ill the assay can be presumed to be accurate. Thus, if the second group does not become ill the test material does not contain prions and if the second group does become ill the test material does contain prions.

By using standardized prion preparations of the invention it is possible to create extremely dilute compositions containing the prions. For example, a composition containing one part per million or less or even one part per billion or less can be created. Such a composition can be used to test the sensitivity of the transgenic mice of the invention in detecting the presence of prions in the sample.

- 51 -

Prion preparations of the present invention are desirable in that they will include a constant amount of prions and are extracted from an isogeneic background. Accordingly, contaminants in the preparations will be  
5 constant and controllable. Standardized prion preparations of the invention will be useful in the carrying out of bioassays in order to determine the presence, if any, of prions in various pharmaceuticals, whole blood, blood fractions, foods, cosmetics, organs  
10 and in particular any material which is derived from an animal (living or dead) such as organs, blood and products thereof derived from living or dead humans. Thus, standardized prion preparations of the invention will be valuable in validating purification protocols  
15 where preparations are spiked and reductions in titer measured for a particular process.

#### Measuring Levels Of Prions

The present invention can be utilized to determine the concentration of prions (which generally only infect  
20 a genetically diverse animal) within a given sample. The transgenic mice make it possible to test for the positive presence of prions within a sample. The mice are capable of detecting the presence of prions in a concentration as low as 1 ppm or even 1 ppb or less. The procedure for  
25 doing such will be apparent to those skilled in the art upon a review of the present disclosure in combination with an article entitled "Measurement Of The Scrapie Agent Using An Incubation Time Interval Assay," published by Prusiner, et al, Annals. of Neurology 11:353-358  
30 (1982) which is incorporated herein by reference to disclose such a method of measurement. In general, the method is carried out by determining the titer of the prions by carrying out measurements of time intervals from inoculation to onset of symptoms and from

- 52 -

inoculation to death. The intervals are inversely proportioned to the size of the dose injected intracerebrally. The logarithms of the time intervals minus a time factor are linear functions of the  
5 logarithms of the inoculum size.

#### Chimeric PrP Gene

Since the fundamental event underlying prion propagation seems to be a conformational change in PrP [Pan et al., Proc. Natl. Acad. Sci. USA 90:10962-10966  
10 (1993)] and mouse PrP differs from human PrP at 31 positions out of 254 [Kretzschmar et al., DNA 5: 15-324 (1986)], we constructed modified PrP transgenes. Chimeric SHa/Mo transgenes have produced prions with new properties, the most useful being the chimeric SHa/Mo  
15 transgene labeled MH2M which carries 5 amino acid substitutions found in SHaPrP lying between codons 94 and 188. [Scott et al., Cell 73:979-988 (1993)]. We made an analogous chimeric human/mouse PrP gene, which we call MHu2M, in which the same region of the mouse gene is  
20 replaced by the corresponding human sequence which differs from mouse PrP at 9 codons as is shown in Figure 2.

Mice expressing the MHu2M chimeric transgene are susceptible to human prions after abbreviated incubation  
25 times. More specifically, the transgenic mice of the present invention which include the chimeric MHu2M gene will, after inoculation with human prions, develop disease symptoms attributed to the prions within about 200 days  $\pm$  50 days. Further, 80% or more the transgenic  
30 mice of the invention inoculated with human prions will develop symptoms of the disease, more preferably 98% or more of the mice will develop symptoms of the disease. According to experiments carried out, 100% of the transgenic MHu2M mice inoculated with human prions

- 53 -

actually developed symptoms of the disease in about 200 days or less  $\pm$  50 days.

These findings indicate that murine cells cannot readily convert HuPrP<sup>C</sup> into HuPrP<sup>Sc</sup> but they can process  
5 MHu2MPrP<sup>C</sup> into MHu2MPrP<sup>Sc</sup>. Since Tg(MHu2M) mice develop neurodegeneration more rapidly than monkeys, they provide a preferred host for bioassays of infectivity in tissues of humans dying of prion diseases. The Tg(MHu2M) mice disclosed herein provide an excellent system for  
10 assessing the sterility of pharmaceuticals as well as tissue and organ grafts prepared from human sources. Other transgenic mice which include the prion protein gene of the animal in danger of infection can be used to test samples for prion diseases which can infect domestic  
15 animals such as sheep and cattle.

Samples for assay may be obtained from any source, including animal and plant sources. Such samples are prepared for inoculation into the transgenic mammal by the methods described herein or methods known to those  
20 skilled in the art.

#### Chimeric MHu2M Gene

Figure 1 is shown regarding how to create the chimeric MHu2M gene. At first, we engineered a new KpnI site in the HuPrP ORF cassette (shown shaded), changing  
25 nucleotide 282 from a cytosine to a thymine residue by PCR-mediated mutagenesis. This mutagenic change conserves the amino acid sequence of HuPrP. A second oligonucleotide primer complimentary to DNA sequences around the BstE11-cut product was used to replace the  
30 corresponding MoPrP gene fragment (the MoPrP gene is unshaded) creating the hybrid gene MHu2M. Microinjection of a cosSHa.Tet construct bearing this expression cassette resulted in founder animal Tg(MHu2M)FVB-B5378.

- 54 -

An expanded representation of the region of MHu2MPPrP between codons 94 and 188 which is flanked by MoPrP sequences (Figure 2). MHu2MPPrP differs from MoPrP by nine amino acids in the region between amino acids 96 and 167. These amino acid residues which are derived from HuPrP are shown on the lower section of the diagram; the amino acids at the same position of MoPrP are shown above. The discrepancy of amino acid positions is due to MoPrP having one less amino acid than HuPrP in the region immediately upstream from the replacement.

#### Artificial PrP Genes

The real power of the present invention lies in the understanding that a variety of different artificial PrP genes can be created which, when inserted into a host animal, will render that animal susceptible to infection with prions which normally only infect a second and genetically diverse test animal. There are nearly an infinite number of possible artificial PrP genes which would meet the basic criteria of the invention, i.e. rendering a mammal such as a mouse susceptible to infection with prions which normally infect only a genetically diverse test animal such as a human. The MHu2M gene of the invention is only one specific example of an artificial gene which achieves the primary object of the invention. By reviewing Figures 3, 4 and 5 numerous other artificial gene possibilities will be deduced by those skilled in the art. Specifically, referring to Figure 3 one can readily determine the amino acid sequence of mouse PrP and observe the positions wherein the mouse PrP sequence differs with a human PrP sequence. Thus, to create an artificial gene one can substitute a codon (or sequence of codons) of a mouse PrP gene with a codon (or sequence of codons) of a human PrP gene at the same position which will encode a different



- 55 -

amino acid - any (but not all) of the codons where different sequences appear can be used for substitution. Mutations in PrP genes can also be taken into consideration and used to create chimeric genes with further advantages as described above. It will be understood by those skilled in the art that, if all of the codons where differences appear between the mouse and the human were substituted, the resulting gene would be the human PrP gene, which is not part of the present invention. However, as explained above, the entire human PrP gene can be inserted into a host animal such as a mouse to create a transgenic animal of the invention, expressing two fold or higher levels of human PrP<sup>C</sup> are included. Transgenic mice expressing only low levels of human PrP<sup>C</sup> are unlikely to become ill after inoculation with human prions. However, if the level of human PrP<sup>C</sup> expression is elevated, the transgenic animals become susceptible to infection with human prions. This is another means of overcoming the species barrier by what is referred to as a stochastic process.

Referring to Figures 1 and 4 it can be seen how it would be possible to produce artificial PrP genes wherein the resulting gene could be inserted into a mouse in order to render the mouse susceptible to infection with bovine prions. A similar strategy with respect to producing a mouse which would be susceptible to infection with sheep prions can be deduced from reviewing Figure 5. In addition to these possibilities those skilled in the art will recognize that, in certain instances, completely artificial nucleotide sequences can be used as corresponding substitutes for a portion of the natural sequence in order to obtain a useful artificial gene which, when inserted into an animal, will render that animal susceptible to infection with prions which normally would infect only a genetically diverse mammal.

- 56 -

Other artificial genes of the invention include genes which may be native or artificial and are operatively fused to an enhanced promoter such as a neuron specific enolase promoter. The enhanced promoter is such that when inserted into the host mammal will express the PrP gene at a level sufficiently high to render the host animal susceptible to infection with prions which normally only infect a genetically diverse test animal.

#### 10 Evidence of Disease

PrP<sup>Sc</sup> has been found in the brains of affected Tg(MHu2M) mice after inoculation with Hu(CJD) or Mo(RML) prions. Brain homogenates of Tg(MHu2M) were either left undigested or digested with proteinase K (BMB) at a final concentration of 20 µg/ml for 1 hour at 37°C (even numbered lanes). Samples were resolved by SDSPAGE and then analyzed by Western blot.

The distribution of PrP<sup>C</sup> and PrP<sup>Sc</sup> in clinically sick Tg(MHu2M) mice inoculated with Mo(RML) and Hu(CJD) prions were detected by the histoblot method. The histoblots included those of coronal sections through the region of the hippocampus and thalamus. Differences are observed between: (A) PrP<sup>C</sup> in Mo(RML) infected mouse; (B) PrP<sup>C</sup> in sporadic CJD RG-infected mouse; (C) PrP<sup>Sc</sup> in Mo(RML) infected mouse; (D) PrP<sup>Sc</sup> in sporadic CJD RG-infected mouse; (E) PrP<sup>Sc</sup> in sporadic CJD EC-infected mouse; and (F) PrP<sup>Sc</sup> in iatrogenic CJD (#364)-infected mouse.

#### EXAMPLES

The following examples are put forth so as to provide those of ordinary skill in the art with a complete disclosure and description of how to make and use the standardized preparation, chimeric genes, transgenic mice and assays of the present invention, and

- 57 -

are not intended to limit the scope of what the inventors regard as their invention. Efforts have been made to ensure accuracy with respect to numbers used (e.g. amounts, temperature, etc.) but some experimental errors and deviations should be accounted for. Unless indicated otherwise, parts are parts by weight, molecular weight is weight average molecular weight, temperature is in degrees Centigrade, and pressure is at or near atmospheric.

10

Example 1Construction of Chimeric Gene (MHu2M)

The source of the HuPrP ORF for construction of an expression cassette has been described [Hsiao et al., Nature 338:342-345 (1989)]. The construction of the MHu2M gene is described in connection with the description of Figure 1. All PrP ORF cassettes were flanked by Sall and XhoI, which cleave immediately adjacent to the PrP initiation and termination codons of the PrP ORF respectively, allowing for convenient subcloning into the cos.SHaTet cosmid expression vector [Scott et al., Cell 73:979-988 (1993)]. The isolation and screening of recombinant cosmid clones was achieved by methods which have been previously described [Scott et al., Cell 73:979-988 (1993)].

25

Example 2Producing Transgenic Mice/Tg(MHu2M)

The nucleotide sequences of the HuPrP and MHu2MPrP ORFs of Example 1 were verified. The cosmid NotI fragments, recovered from large-scale DNA preparations, were used for microinjection into the pronuclei of fertilized C57BL/6 X SJL or FVB/N oocytes as previously described [Scott et al., Cell 59:847-857 (1989); Scott et al., Protein Sci. 1:986-997 (1992)]. Genomic DNA

- 58 -

isolated from tail tissue of weaning animals was screened for the presence of incorporated transgenes using a probe that hybridizes to the 3'-untranslated region of the SHaPrP gene contained in the cosSHa.Tet vector [Scott et al., Protein Sci. 1:986-997 (1992)]. The offspring obtained were tested and it was confirmed that the chimeric MHu2M gene was integrated into the genome of these offspring. As shown in Example 5 below, these mice were found to be susceptible to infection with human prions 100% of the time.

### Example 3

#### Preparation of Brain Homogenates

A 10% (w/v) homogenate of a sample of thawed human brain tissue was prepared in phosphate buffered saline lacking calcium and magnesium ions. The tissue was initially dissociated using a sterile disposable homogenizer, and this suspension was subjected to repeated extrusion through an 18 gauge syringe needle followed by a 22 gauge needle. Samples for inoculation into test animals were diluted 10-fold. Homogenates of clinically sick Tg and non-Tg mouse brains were prepared in the same way except for the omission of the initial dissociation step.

- 59 -

Example 4Sources of Prion Inocula

Human inocula were derived from frozen brain tissues of patients in which the clinical diagnosis of CJD or GSS  
5 had been confirmed by histopathological examination of brain tissues and, in most cases, by prion protein analysis. In some cases, the PrP gene was amplified by PCR of DNA isolated from patient blood and the PrP sequence determined by DNA sequence analysis. No HuPrP  
10 mutations were detected in cases of sporadic or iatrogenic CJD. The RML isolate was obtained from Swiss mice [Chandler, R.L., Lancet 1:1378-1379 (1961)] from a closed colony at the Rocky Mountain Laboratory or in Swiss CD-1 mice obtained from Charles River Laboratories.

15

Example 5Determination of Scrapie Incubation Periods

Transgenic mice as per Example 2 were inoculated intracerebrally with 30  $\mu$ l of brain extract using a 27 gauge needle inserted into the right parietal lobe.  
20 The preparation of inocula and criteria for diagnosis of scrapie in mice have been described [Carlson et al., Cell 46:503-511 (1986)]. Beginning 50 days after inoculation, the mice were examined for neurologic dysfunction every 3 days. When clinical signs of scrapie appeared, the mice  
25 were examined daily. When some animals whose death was obviously imminent were identified, their brains were taken for histopathologic studies (as per the procedures of Example 3) and confirmation of the diagnosis of scrapie.

- 60 -

Example 6Immunoblot Analysis

For the determination of the relative levels of PrP expression in Tg mouse and human brains, protein concentrations were determined by bicinchoninic acid assay and immuno dots blots as previously described [Scott et al., Cell 73:979-988 (1993)]. Samples for Western blot analysis were prepared and western blots were performed as described previously [Towbin et al., 10 Proc. Natl. Acad. Sci. USA 76:4350-4354 (1979)], except that an enhanced chemiluminescent (ECL) detection method (Amersham) was used. The blot was exposed to X-ray film for 5-60 seconds.  $\alpha$ -PrP R073 rabbit antiserum was used at a final dilution of 1:5000 and 3F4 monoclonal antibody 15 was also employed [Serban et al., Neurology 40:110-117 (1990)].

Example 7Tg(MHu2MPrP) Mice Are Susceptible to Human Prions

Using information from a systematic study of 20 chimeric SHa/Mo PrP genes, a chimeric Hu/Mo PrP gene analogous to MH2M was constructed as per Example 1. This gene, which we call MHu2M, differs from MoPrP at 9 positions, all of which lie between codons 94 and 188 as is shown in Figure 2. A Tg mouse was constructed 25 expressing the MHu2M PrP gene and the founder designated Tg(MHu2M)FVB-B5378 as per Example 2. By serial dilution of Tg(MHu2M) mouse and human brain homogenates, we estimate that the level of MHu2MPrP<sup>C</sup> in the brains of these Tg(MHu2M)FVB-B5378 mice are similar to those of 30 HuPrP<sup>C</sup> found in human brain using an immuno dot blotting procedure.

Tg(MHu2M) mice as per Example 2 were inoculated with brain homogenates from three unrelated Caucasian patients who had died of CJD. Two of the three patients died of



- 51 -

sporadic CJD: one (R.G.) was a 56-year-old American female; the other (E.C.) was a 61-year-old Canadian female. In both cases, cerebrocortical biopsy showed severe spongiform degeneration. The third (#364) was a British youth who had contracted iatrogenic CJD after treatment for hypopituitarism with human growth hormone (HGH) derived from cadaveric pituitaries [Collinge et al., Lancet 337:1441-1442 (1991)]. Brain homogenates from all three CJD patients exhibited protease-resistant PrP by Western immunoblotting. This protease-resistant isoform of PrP is designated PrP<sup>Sc</sup>, or often PrP<sup>CJD</sup> when it is found in humans.

All of the Tg(MHu2M) mice inoculated with homogenates from the CJD patients described above developed signs of central nervous system (CNS) dysfunction ~200 days after inoculation (see Table 1 below). The clinical signs were similar to those of murine scrapie. After developing clinical signs, the inoculated Tg(MHu2M) mice died rapidly, often within a day. Two of the eight uninoculated Tg(MHu2M) mice are now >500 days of age and remain well; the youngest of the other six uninoculated animals is older than the age at which inoculated Tg(MHu2M) mice developed signs of CNS dysfunction.

Inoculation of Tg(MHu2M) mice with Mo(RML) prions passaged in mice produced disease in 178 ± 3 days, which is ~40 longer than Mo(RML) prions in non-Tg mice. Prolongation of incubation times in mice expressing non-murine transgenes is well established, and occurs presumably because the foreign PrP<sup>C</sup> molecule inhibits the conversion of MoPrP<sup>C</sup> into MoPrP<sup>Sc</sup> [Prusiner et al., Cell 63:673-686 (1990)]. In contrast to Tg(MHu2M) mice, incubation times for RML prions in Tg(MH2M) mice were the same as those of the non-Tg mice [Scott et al., Cell 73:979-988 (1993)].

- 62 -

Table 1. Incubation of human (CJD) and mouse (RML) prion inocula in Tg(MHu2M)FVB-B5J/8 mice

5	Source	Inoculum	No. <sup>a</sup>	Incubation Times (mean days $\pm$ SE)		Death <sup>b</sup>
				Range (days)	Illness	
	Sporadic CJD	RG	8/8	225-249	238 $\pm$ 3.2	240 $\pm$ 5.4 (3)
	Sporadic CJD	EC	7/7	202-229	218 $\pm$ 4.6	N.D.
10	Iatrogenic CJD	#364	9/9	221-245	232 $\pm$ 2.9	235 $\pm$ 3.9 (5)
	Mo	RML	19/19	155-195	178 $\pm$ 3.3	203 $\pm$ 2.0 (14)

<sup>a</sup> Number of animals developing clinical sickness (neurological dysfunction) divided by the total number of animals inoculated. In the case of inoculum RG, three animals were found dead after 224, 238, and 243 days before a clinical diagnosis could be made. In the case of inoculum EC, two animals were found dead after 225 and 226 days before a clinical diagnosis could be made. In each case, these animals died at the same time that clinical diagnosis was made in other inoculated animals.

<sup>b</sup> The number of mice dying of scrapie is shown in parentheses. Mice sacrificed for pathological examination are excluded from these calculations.

25

Example 8Comparative ExampleTg(HuPrP) Mice Are Resistant to Human Prions

Tg mice expressing HuPrP were produced using the HuPrP gene ORF, which had been cloned into the cosSha.Tet expression vector [Scott et al., Protein Sci. 1:986-997 (1992)]. Microinjection of outbred C57B6/SJL and inbred FVB mouse embryos resulted in two founder transgenic animals designated Tg(HuPrP)B6SJL-110 and Tg(HuPrP)FVB-152. We estimated by serial dilution of

- 63 -

brain homogenates and immuno dot blotting, that the level of PrP<sup>c</sup> in the brains of the progeny of these founders express HuPrP at levels 4- to 8-fold higher than the level of HuPrP found in the human brain.

5 To determine whether expression of HuPrP in Tg(HuPrP)B6SJL-110 and Tg(HuPrP)FVB-152 conferred susceptibility to human prions, incubation periods were measured after inoculation of Tg(HuPrP) and non-Tg mice with brain extracts from 18 patients that had died of  
10 sporadic CJD, iatrogenic CJD, familial CJD or GSS. From experiments performed over the past 2.5 years, we concluded that the two lines of Tg(HuPrP) mice were no more responsive than non-Tg mice to human prions (see Table 2 below). The rate of transmission to Tg(HuPrP)  
15 mice was 8.3% (14 clinically sick mice out of 169 mice) which was similar to a transmission rate of 10.3% in control non-Tg mice (6 clinically sick mice out of 58 mice). In the infrequent event of a positive transmission, incubation times were extremely long  
20 ranging, from 590 days to 840 days in both Tg(HuPrP) and non-Tg mice. By this late time, many animals had died of intercurrent illnesses which complicated diagnosis. The difficulty of interpreting transmissions occurring after extremely long incubation periods is compounded by the  
25 heightened potential for artifactual results due to low levels of contaminating prions.

Statistical analysis shows that the frequency of Hu prion transmission to Tg(MHu2MPrP) mice compared to Tg(HuPrP) and non-Tg mice is highly significant using the  
30 Fisher's exact test,  $p < 10^{-7}$  [Mehta et al., J. Am. Stat. Assn. 78:(392) 427-434 (1983)]. When Hu prion transmission to Tg(HuPrP) mice was compared to non-Tg mice, the frequencies were similar,  $p = 0.79$ .

To confirm the clinical diagnosis of prion disease,  
35 5 ill Tg(HuPrP) and 1 non-Tg mice were sacrificed and

- 64 -

brain extracts were examined for the presence of PrP<sup>Sc</sup> by Western blotting with the  $\alpha$ -PrP antibodies, 3F4 mAb and R073 antiserum [Kascsak et al., J. Virol. 61:3688-3693 (1987); Serban et al., Neurology 40:110-117 (1990)]. The

5 3F4 mAb reacts specifically with HuPrP allowing discrimination from MoPrP. MoPrP<sup>Sc</sup> was detected in the brain of the non-Tg mouse inoculated with sporadic CJD inoculum #87011 which developed clinical signs after 756 days, while 3F4-reactive PrP<sup>Sc</sup> was detected in the brains

10 of two Tg(HuPrP) mice which developed clinical signs after 589 days post-inoculation with iatrogenic CJD inoculum #170. The equivalent transmission rates of human prions in Tg(HuPrP) and non-Tg mice indicate that this is a rare event with the same frequency of

15 occurrence as the stochastic conversion of MoPrP<sup>C</sup> to MoPrP<sup>Sc</sup> induced by human prions.

The absence of either R073- or 3F4-reactive PrP<sup>Sc</sup> in the brains of 3 out of the 6 mice analyzed may reflect the difficulty of accurately diagnosing prion disease in

20 elderly animals. Some of the mice inherited prion diseases of both humans and Tg mice exhibit little or undetectable levels of protease-resistant PrP; yet, based on transmission studies, their brains contain prions and they show clear spongiform degeneration [Medori et al.,

25 N. Engl. J. Med. 326:444-449 (1992)].

In contrast to Tg(MHu2M) mice, Hu prions from patient RG have not transmitted to either Tg(HuPrP) or non-Tg mice after >330 days (see Table 2 below).

Attempts to transmit preparations enriched for Hu prion

30 rods prepared from the brain of patient RG have likewise been negative for >300 days. In addition, inoculum from the iatrogenic CJD case (#364) has produced illness in neither Tg(HuPrP) nor non-Tg mice after >780 days (as shown in Table 2 below).

TABLE 2

Table 2. Incubation times in Tg(HuPrP)FVB-152 and Tg(HuPrP)E6SJL-110 mice after inoculation with brain extracts from patients with human prion diseases

	Host	Inoculum	(n/n <sub>o</sub> )	Incubation times (days $\pm$ SE) <sup>b</sup>
5	Tg152	Sporadic CJD(#87011)	1/10	706
	Non-Tg	Sporadic CJD(#87011)	3/5	697.3 $\pm$ 51
	Tg 152	Sporadic CJD(#88037)	3/10	680 $\pm$ 28
10	Tg 152	Sporadic CJD(RG)	0/10	
	Non-Tg	Sporadic CJD(RG)	0/10	
	Tg 152	Sporadic (RG) Rods	0/8	
	Non-Tg	Sporadic (RG) Rods	0/8	
	Tg 152	Codon 102 GSS(#87027)	4/10	724 $\pm$ 16
15	Non-Tg	Codon 102 GSS(#87027)	0/10	679
	Tg 152	Codon 102 GSS(#87031)	0/10	
	Non-Tg	Codon 102 GSS(#87031)	1/5	742
	Tg 152	Codon 178 F-CJD	0/8	
	Non-Tg	Codon 178 F-CJD	0/8	
20	Tg 110	Sporadic CJD(#87036)	0/8	
	Non-Tg	Sporadic CJD(#87036)	1/5	838
	Tg 110	Iatrogenic CJD(#703)	0/10	
	Non-Tg	Iatrogenic CJD(#703)	0/5	
	Tg 110	Iatrogenic CJD(#170)	2/10	589 $\pm$ 0

- 66 -

	Non-Tg	Iatrogenic CJD(#170)	0/5	
	Tg 110	Iatrogenic CJD(#364)	0/10	
	Non-Tg	Iatrogenic CJD(#364)	0/5	
	Tg 110	Codon 200 F-CJD	1/8	791
5	Tg 110	Codon 217 GSS	1/8	874
	Tg 110	Codon 102 GSS-A	0/10	
	Tg 110	Codon 102 GSS-B	1/8	694
	Tg 110	Codon 117 GSS	0/8	

\* Number of animals developing clinical sickness divided by the total number of animals inoculated.

" Refers to time to diagnosis of illness.

Patients from which inoculum were derived are described in the following publications: [Collinge et al., Lancet 337:1441-1442 (1991); Hsiao et al., Nature 338:342-345 (1989); Hsiao et al., Neurology 41:681-684 (1991)].

#### Example 9

##### Formation of MHu2MPrP<sup>Sc</sup> or MoPrP<sup>Sc</sup>

##### In the Brains of Tg(MHu2M) Mice

20 Some clinically sick Tg(MHu2M) mice inoculated with each of the three CJD prion inocula or RML prions were sacrificed for histopathological verification of disease and for prion protein analysis. Western blots of brain homogenates from Tg(MHu2M) mice infected with Hu prions  
25 probed with RO73 and 3F4  $\alpha$ -PrP antibodies revealed the presence of protease-resistant PrP<sup>Sc</sup> which reacted with the 3F4 monoclonal antibody showing this protease-resistant product to be MHu2M PrP<sup>Sc</sup>. The epitope



- 67 -

recognized by this antibody consists of a pair of methionine residues at positions 109 and 112 in PrP [Rogers et al., J. Immunol. 147:3568-3574 (1991)] which are contained in MHu2M but not in MoPrP as can be seen by  
5 the mouse/human comparison of figure 2. The polyclonal rabbit  $\alpha$ -PrP antiserum R073 diluted 1:5000 was poorly reactive with MHu2MPrP<sup>Sc</sup> as well as HuPrP<sup>C</sup> and HuPrP<sup>Sc</sup> from diseased human brains. Brain homogenates from Tg(MHu2M) mice infected with RML prions contained PrP<sup>Sc</sup> which was  
10 detectable only with R073 and not 3F4  $\alpha$ -PrP antibodies, indicating that Tg(MHu2M) mice are capable of producing MoPrP<sup>Sc</sup> but not MHu2MPrP<sup>Sc</sup> in response to RML prions previously passaged in mice. While these findings are similar to those reported for Tg(SHaPrP) mice [Scott  
15 et al., Cell 59:847-857 (1989)], they contrast with those found for Tg(MH2MPrP) mice where MH2MPrP<sup>Sc</sup> was formed in response to RML prions [Scott et al., Cell 73:979-988 (1993)].

#### Example 10

#### Regional Distribution of PrP<sup>Sc</sup> and Patterns of Neuropathology

The distribution of Mo and MHu2M PrP<sup>C</sup> and PrP<sup>Sc</sup> is shown in histoblots of coronal brain sections through the hippocampus and thalamus of Tg(MHu2M) mice inoculated  
25 with RML or CJD prions. The weak immunoreactivity of MHu2M PrP with R073 permitted a degree of analysis which had not been previously possible in Tg mice expressing SHaPrP or MH2MPrP because these PrP species react with this antibody. The pattern of PrP<sup>Sc</sup> deposition was highly  
30 dependent upon the species of origin of the infectious prions. When inoculated with RML prions, histoblots of the brains of Tg(MHu2M) were similar to those of CD-1 mice infected with RML prions, revealing a diffuse pattern of MoPrP<sup>Sc</sup> deposition in the hippocampus,

- 68 -

thalamus, hypothalamus and all layers of the neocortex. The histoblot pattern of was strikingly different for Tg(MHu2M) mice inoculated with CJD prions. The deposition of MHu2MPrP<sup>Sc</sup> was confined primarily to the  
5 deep layers of the neocortex, the thalamus, particularly the ventral posterior medial thalamic nucleus, the hypothalamus and the putamen. The hippocampal region and the outer layers of the neocortex were spared. Interestingly, while the hippocampus was completely  
10 devoid of MHu2MPrP<sup>Sc</sup>, this region showed the most intense MHu2MPrP<sup>C</sup> signal. The same pattern of MHu2MPrP<sup>Sc</sup> deposition was consistently observed in histoblots of Tg(MHu2M) mice inoculated with all three CJD prion isolates prepared from human brain. It is noteworthy  
15 that the pattern of MHu2MPrP<sup>Sc</sup> deposition is similar to the pattern of PrP<sup>CJD</sup> deposition observed in histoblots of the brain from which inoculum RG was derived [Taraboulos et al., Proc. Natl. Acad. Sci. USA 89:7620-7624 (1992)]. The spongiform degeneration in the brains of Tg(MHu2M)  
20 mice infected with Mo(RML) and Hu(CJD) prions reflected the patterns of PrP<sup>Sc</sup> accumulation described above.

#### Experimental Protocols

Numerous additional examples of transgenic and hybrid mice as well as comparative examples and methods  
25 of testing such are described below. These examples and methods are listed in Tables 3-7. With respect to such the (1) methods of making mice; (2) brain homogenates; (3) prion inocula; (4) measurement of incubation times; (5) immunoblotting; and (6) immunohistochemistry are  
30 described below.

- 69 -

Production Of Transgenic Mice

The MoPrP-A sequence used was derived from Swiss CD-1 mice Locht, C., et al., "Molecular cloning and complete sequence of prior protein cDNA from mouse brain infected with the scrapie agent," Proc. Natl. Acad. Sci. USA 83:6372-6376 (1986). Construction of the MoPrP ORF cassette has been previously described Rogers, M., Serban, D., Gyuris, T., Scott, M., Torchia, T., and Prusiner, S.B. (1991). Epitope mapping of the Syrian hamster prior protein utilizing chimeric and mutant genes in a vaccinia virus expressions system. J. Immunol. 147:3568-3574. The construction of the MoPrP-P101L expression cassette containing a substitution of a leucine for proline codon at residue 101 of the mouse PrP gene, corresponding to the GSS mutation at codon 102 of the human PrP gene has been described Hsiao, K. and Prusiner, S.B. (1990). Inherited human prion diseases. Neurology 40:1820-1827. ORF cassettes were digested with BglII (which cleaves immediately adjacent to the initiation codon). The 5' protruding termini were filled in using Klenow, and SalI linkers were added. Recombinant clones were cleaved with SalI and XhoI (which cleaves immediately past the PrP termination codon), and purified ORF fragments were ligated to the SalI-cut cosSHA.Tet cosmid expression vector Scott, M.R., Köhler, R., Foster, D., and Prusiner, S.B. (1992). Chimeric prion protein expression in cultured cells and transgenic mice. Protein Sci. 1:986-997. The isolation of recombinant cosmid clones was achieved by methods that have been previously described Scott, M., Groth D., Foster, D., Torchia, M., Yang, S.-L., DeArmond, S.J., and Prusiner, S.B. (1993). Propagation of prions with artificial properties in transgenic mice expressing chimeric PrP genes. Cell 73:979-988. NotI fragments, recovered from large-scale DNA cosmid preparations, were

- 70 -

- used for microinjection into the pronuclei of fertilized FVB/N oocytes as previously described Scott, M., Foster, D., Mirenda, C., Serban D., Coufal, F., Wälchli, M., Growth, D., Carlson, G., DeArmond, S.J.,
- 5 Westaway, D., and Prusiner, S.B. (1989). Transgenic mice expressing hamster prion protein produce species-specific infectivity and amyloid plaques. Cell 59:847-857. Genomic DNA isolated from tail tissue of weaning animals was screened for the presence of
- 10 incorporated transgenes using a probe that hybridizes to the 3'-untranslated region of the SHaPrP gene contained in the cosDHa.Tet vector Scott, M.R., Köhler, R., Foster, D., and Prusiner, S.B. (1992).

#### Preparation Of Brain Homogenates

- 15 Ten % [w/v] homogenates of mouse brain were prepared by repeated extrusion through an 18 gauge syringe needle followed by a 22 gauge needle in phosphate buffered saline (PBS) lacking calcium and magnesium ions. For immunoblot analysis, samples were cleared of cell debris
- 20 by a brief low-speed centrifugation. Purified Hu prions were prepared using a protocol previously developed for SHa prions Prusiner et al., (1983) Scrapie Prions Aggregate to Form Amyloid-like Birefringent Rods. Cell 35, 349-358.

#### 25 Prion Inocula

- Human brain specimens were collected from patients dying of sporadic, inherited or infectious prion disease. The iatrogenic CJD denoted 364 was provided by Dr. John Collinge. The RML isolate from Swiss mice Chandler,
- 30 R.L., "Encephalopathy in mice produced by inoculation with scrapie brain material," Lancet 1:1378-1379 (1961) was provided by Dr. William Hadlow and was passaged in Swiss mice CD-1 mice obtained from Charles River

- 71 -

Laboratories (Wilmington, MA). For transmission of neurodegeneration from spontaneously ill Tg(MoPrP-P101L) mice, brain homogenates were prepared.

#### Measurement Of Incubation Times

- 5 Mice were inoculated intracerebrally with 30  $\mu$ l of samples prepared from brain using 27 gauge needle inserted into the right parietal lobe, and observed for signs of disease. Samples were diluted 10-fold in PBS prior to inoculation. Additionally, uninoculated
- 10 Tg(MoPrP-P101L) and Tg(MoPrP-A) mice were observed for spontaneous CNS disease. The preparation of inocula and criteria for diagnosis of scrapie in mice have been described Carlson, G.A., et al., "Lineage of prion protein and scrapie incubation time genes," Cell 46:503-511
- 15 (1986). When clinical signs of CNS dysfunction appeared, the mice were examined daily. To confirm the clinical diagnosis, the brains of some animals whose deaths were obviously imminent were taken for histopathological studies.

#### 20 Immunoblotting

- Total protein concentrations in brain homogenates were determined by bicinchoninic acid assay. Immuno dot blots for the determination of the relative levels of PrP expression in Tg mouse brains were performed as
- 25 previously described (Scott et al., 1993). Samples for Western blot analyses were prepared by digesting brain homogenates with 20  $\mu$ g proteinase K for 60 min at 37°C. Western blots were performed as described previously Barry, R.A., et al., "Monoclonal antibodies to the
- 30 cellular and scrapie prion proteins," J. Infect. Dis., 154:518-521 (1986); Towbin, H., et al., "Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose sheets: Procedure and some applications,"

- 72 -

Proc. Natl. Acad. Sci. USA 76:4350-4354 (1979), except that an enhanced chemiluminescent (ECL) detection method (Amersham, Arlington Heights, IL) was used. The lot was exposed to X-ray film  $\alpha$ -PrP R073 rabbit antiserum was used at a final dilution of 1,5000.

#### Immunohistochemistry

For immunohistochemistry, endogenous peroxidase activity was blocked with methanol-hydrogen peroxide (2 ml of 30%  $H_2O_2$  in 200 ml methanol). To enhance PrP immunoreactivity, the sections were immersed in 1.3 mM HCl and autoclaved at 121°C for 10 min Muramoto et al., (1992) The sequential development of abnormal prion protein accumulation in mice with Creutzfeldt-Jakob disease. Am. J. Pathol. 140, 1411-1420. When temperature decreased, the slides were placed under running tap water for 10 min followed by rinsing with TB (50 mM Tris-HCl, pH 7.6, with 150 mM NaCl). Nonspecific protein binding was blocked with TBS (25 mM Tris-HCl, pH 7.8; 0.05% Tween 20; 0.5 M NaCl; and 5% nonfat milk) for 10 min. The  $\alpha$ -PrP polyclonal rabbit R073 antiserum was diluted 1:250 in TBS. Tissue sections were incubated with the primary antiserum overnight at 4°C. Following 2 rinses with TB containing 0.1% Triton-X and once in TB, the sections were incubated with biotinylated goat anti-rabbit IgG (Vector Laboratories, Burlingame, CA) in TBS, 1:100 dilution for 30 min at room temperature. After 2 rinses with TB containing 0.1% Triton-X, the sections were incubated with an avidin-biotin-peroxidase complex (Vector Laboratories) for 30 min at room temperature. Three rinses in TB were then followed by a 3-5 min reaction with diaminobenzadine solution (5 mg diaminobenzadine tetrahydrochloride, 68 mg imidazole, and 7 mg NaN<sub>3</sub> in 10 ml of TB), to which 100  $\mu$ l of 0.0015%  $H_2O_2$  was added.



- 73 -

Peroxidase immunohistochemistry with antibodies to glial fibrillary acidic protein was used to evaluate the extent of reactive astrocytic gliosis. Hydrolytic autoclaving was used to detect PrP immunoreactivity  
5 (Muramoto et al., 1992).

#### EXAMPLES RE TABLES 3-7

##### Transgenic Mice With Human and Chimeric PrP Genes

FVB mice expressing human, chimeric Hu/Mo and mutant PrP genes were constructed using the cos.SHaTet cosmid  
10 expression vector derived from the Syrian hamster (SHa) (Scott et al., 1992). Table 3 below shows the designation of the mouse line, the expressed PrP<sup>C</sup> molecules and the approximate level of transgenic expression. Also indicated are those mouse lines that  
15 were crossed with Prnp<sup>0/0</sup> mice in which the mouse PrP gene had been disrupted by homologous recombination (Büeler et al., 1992). Backcrossing these mice produced animals in those encoded by the transgene. While SV129ES cells were used to generate a chimeric mouse with a disrupted PrP  
20 allele, that mouse was mated with a C57BL mouse and the offspring crossed to each other to produce null animals. Subsequently, these Prnp<sup>0/0</sup> mice were repeatedly crossed onto the FVB background.

TABLE 3

Nomenclature and Characteristics of Transgenic Mouse Lines				
	Mouse Line Description	Expressed PrP <sup>c</sup> Molecules	PrP Transgene Expression	Sequence <sup>b</sup>
5	(i) Tg (HuPrP) mice			
	Tg (HuPrP) 152/FVB	Hu, Mo	~4-8x	V129
10	Tg (HuPrP) 152/Prnp <sup>0/0</sup>	Hu	~4-8x	V129
	Tg (HuPrP) 440/Prnp <sup>0/0</sup>	Hu	~2x	M129
15	(ii) Tg (MHu2M) mice			
	Tg (MHu2M) 5378/FVB	MHu2M, Mo	~1x	M128
	Tg (MHu2M) 5378/Prnp <sup>0/0</sup>	MHu2M	~1x	M128
20	Tg (MHu2M-P101L) 69/Prnp <sup>0/0</sup>	MHu2M-L	~2x	M128, L10

Since the Hu prion inocula are brain homogenates or  
 25 purified prion rods from a variety of patients who died  
 of prion disease, the designations for the patients as  
 well as clinical phenotypes are listed in Table 4 below.  
 The PrP genotypes of the patients are also described.

- 75 -

TABLE 4

Brain Inocula From Patients Who Died of Prion DiseaseSporadic Inocula and Infectious CJD prions  
Containing wt PrP<sup>Sc</sup>

5	Human Inoculum	Prion Disease	Genotype of PrP <sup>d</sup>
	PG	sporadic CJD	wt, M/M129
	EC	sporadic CJD	wt, M/M129
	MA	sporadic CJD	wt, M/M129
	PO	sporadic CJD	wt, M/M129
10	PC	sporadic CJD	wt, M/M129
	364	iatrogenic CJD	wt, M/M129

GSS and Familial CJD prions containing mutant PrP<sup>Sc</sup>

	<u>JJ</u>	<u>GSS</u>	<u>P102L, V/V128</u>
	LJ-1	familial CJD	E200K, M/M129
15	CA	familial CJD	E200K, M/M129
	FH	familial CJD	E200K, V/M129

\*Substitution of L for P codon 102 in HuPrP or at codon 101 in MoPrP or chimeric MHu2MPrP is denoted as "-L".

<sup>b</sup>Level of PrP transgene expression in brain was measured  
20 by serial dilution of the samples followed by dot  
immunoblotting. Each sample was compared to PrP<sup>c</sup> in  
non-Tg mouse or human brain.

<sup>c</sup>Amino acid residues at codon 129 or codon 101.

<sup>d</sup>Patients with sporadic or iatrogenic CJD had wild-type  
25 (wt) PrP ORFs. The PrP alleles encode either M or V at  
position as noted. Mutations in the PrP gene are denoted  
by the wt amino acid followed by the codon number and the  
mutant residue.

- 76 -

MoPrP<sup>c</sup> Inhibits Propagation of Human Prions in Tg(HuPrP) Mice

When Tg(HuPrP)152/FVB mice and non-Tg littermates were inoculated with Hu prions from sporadic or  
5 iatrogenic CJD as well as inherited prion disease cases, ~10% of each group of mice developed CNS dysfunction (Telling et al., 1994). Some of the ill mice produced MoPrP<sup>sc</sup> and others HuPrP<sup>sc</sup> based on Western immunoblots developed with polyclonal  $\alpha$ -PrP antiserum that reacts  
10 with both Hu and MoPrP and with  $\alpha$ -PrP monoclonal antibodies (mAb) that react with Hu but not MoPrP. Those mice that produced HuPrP<sup>sc</sup> demonstrated that HuPrP<sup>sc</sup> could be formed in Mo cells but the process was too infrequent for further study.

15 After Crossing the Tg(HuPrP) 152/FVB mice onto the Prnp<sup>0/0</sup> background, they became susceptible to Hu prions (Table 5)

When Tg(HuPrP)152/FVB mice were inoculated with Hu prions from a case of sporadic CJD, referred to as RG,  
20 only one Tg mouse out of a group of 10 developed neurologic symptoms at >720 d, while non-Tg littermates responded similarly with one animal out of a group of 10 inoculate mice developing neurologic symptoms at >700 d.

- 77 -

TABLE 5

Transmission Of Hu Prions to Tg(HuPrP)/Prnp<sup>0/0</sup> mice

	Recipient Mouse Line	Inoculum <sup>a</sup>	Incubation Time mean d ± SEM (n/no)
5	(A) Tg(HuPrP) FVB Mice		
	Tg(HuPrP) 152/FVB	sCJD(RG)	721 ± 0 (1/10) <sup>†</sup>
	Non-Tg 152/FVB	sCJD(RG)	701 ± 0 (1/10) <sup>b</sup>
	Tg(HuPrP) 152/FVB	sCJD(RG, purified rods)	677 ± 0 (1/10)
	Non-Tg 152/FVB	sCJD(RG, purified rods)	643 ± 42 (3/10)
10	(B) Tg(HuPrP) Prnp <sup>0/0</sup> mice		
	Tg(HuPrP) 152/Prnp <sup>0/0</sup>	sCJD(RG)	263 ± 2 (6/6)
	Tg(HuPrP) 152/Prnp <sup>0/0</sup>	sCJD(EC)	254 ± 6 (9/9)
	Tg(HuPrP) 152/Prnp <sup>0/0</sup>	iCJD(364)	262 ± 8 (5/5)
	Tg(HuPrP) 440/Prnp <sup>0/0</sup>	iCJD(364)	164 ± 2 (7/7)

<sup>a</sup>All samples were 10% (w/w) brain homogenates unless otherwise noted that were diluted 1:10 prior to inoculation. sCJD is sporadic CJD and iCJD is iatrogenic CJD. Patients initials referring to inocula in Table 1B are given in parenthesis.

<sup>b</sup>Transmissions previously reported (Telling et al., 1994).

- 78 -

Similar rates of transmission were observed when Tg(HuPrP)152/FVB and non-Tg mice were inoculated with a preparation highly enriched for PrP<sup>Sc</sup> prepared from the brain of RG (see Section B of Table 5). Using the  $\alpha$ -PrP 3F4 monoclonal antibody (mAb) Kascsak, R.J., et al., "Mouse polyclonal and monoclonal antibody to scrapie-associated fibril proteins," J. Virol. 61:3688-3693 (1987), we estimated, by serial dilution and dot immunoblotting of brain homogenates which were normalized for protein concentration, the level of HuPrP<sup>C</sup> in brains of the Tg(HuPrP)152/FVB mice to be ~4-8 fold higher than HuPrP<sup>C</sup> in human brain (Table 3).

Since earlier studies had shown that heterologous PrP<sup>C</sup> inhibited the conversion of PrP<sup>Sc</sup> as manifest by prolongation of the incubation time Büeler, H., et al., "Mice devoid of PrP are resistant to scrapie," Cell 73:1339-1347 (1993); Prusiner, S.B., et al., "Immunologic and molecular biological studies of prion proteins in bovine spongiform encephalopathy," J. Infect. Dis. 167:602-613 (1993); Prusiner, S.B., et al., "Transgenic studies implicate interactions between homologous PrP isoforms in scrapie prion replication," Cell 63:673-686 (1990), we removed MoPrP<sup>C</sup> by producing Tg(HuPrP)152/Prnp<sup>0/0</sup> mice. When Tg(HuPrP)152/Prnp<sup>0/0</sup> were inoculated with Hu prions, they developed signs of neurologic dysfunction with incubation times between 260 and 300 d (Table 5 shown in Section B).

#### MoPrP Gene Ablation In Mice Expressing Chimeric PrP

Crossing the transgene array from the already susceptible Tg(MHu2M)5378/FVB mice onto the Prnp<sup>0/0</sup> background resulted in a decrease (~20%) in CJD incubation times (Table 6 Sections A and B). Using the  $\alpha$ -PrP 3F4 mAb, we estimated the level of MHu2MPrP<sup>C</sup> in the



- 79 -

brains of the Tg(MHu2m)5378/FVB mice to be slightly less than HuPrP<sup>C</sup> in human brain.

TABLE 6

Transmission of Hu prions to Tg(MHu2MPrP)mice

5	<u>Inoculum*</u>	<u>Incubation Time</u>
		mean d $\pm$ SEM (n/no)
	(A) Tg(MHu2M)/FVB mice inoculated with sporadic or infectious CJD	
	sCJD(RG)	238 $\pm$ 3 (8/8) <sup>b</sup>
	sCJD(EC)	218 $\pm$ 5 (7/7) <sup>t</sup>
10	iCJD(364)	232 $\pm$ 3 (9/9) <sup>o</sup>
	iCJD(364) <sup>c</sup>	231 $\pm$ 6 (9/9)
	sCJD(MA)	222 $\pm$ 1 (4/4)
	(B) Tg(MHu2M)/Prnp <sup>0/0</sup> mice inoculated with sporadic or infectious CJD	
15	sCJD(RC)	202 $\pm$ 2 (6/10)
	sCJD(RG)	191 $\pm$ 3 (10/10)
	iCJD(364)	192 $\pm$ 6 (8/9)
	iCJD(364) <sup>c</sup>	190 $\pm$ 6 (8/8)
	sCJD(MA)	180 $\pm$ 5 (8/8)
20	sCJD(RO)	217 $\pm$ 2 (9/9)
	(C) Tg(MHu2M)/Prnp <sup>0/0</sup> mice inoculated with inherited GSS or CJD	
	GSS(JJ, P102L)	>280 (0/10)
	fCJD(LJ1, E200K)	170 $\pm$ 2 (10/10)
25	fCJD(CA, E200K)	180 $\pm$ 9 (9/9)
	fCJD(FH, E200K)	>250 (0/7)

- 80 -

- \*All samples were 10% (w/v) brain homogenates unless otherwise noted that were diluted 1:10 prior to inoculation. sCJD is sporadic CJD, iCJD is iatrogenic CJD, GSS is Gerstmann-Straussler-Scheinker disease with the codon 102 mutation and fCJD is familial CJD with the codon 200 mutation. Patients initials referring to inocula in Table 4 are given in parenthesis. If the PrP gene of the patient carried a mutation, then the mutation is noted after the patients initials.
- 10 \*Transmissions previously reported (Telling et al., 1994).  
\*This is a second inoculum prepared from a different brain region of iatrogenic CJD patient 364.

Any comparison between the incubation times of Tg(HuPrP)152/Prnp<sup>0/0</sup> and Tg(MHu2M)5378/Prnp<sup>0/0</sup> mice must take into account the different levels of transgene expression. Generally, the level of transgene expression is inversely related to the length of the incubation time. Although the incubation times are similar for Tg(HuPrP)152/Prnp<sup>0/0</sup> and Tg(MHu2M)5378/Prnp<sup>0/0</sup> mice inoculated with Hu prions (Tables 5 and 6 Section B of each), the Tg(HuPrP)152/Prnp<sup>0/0</sup> express 5-10-fold more of the transgene product than Tg(MHu2M)5378/Prnp<sup>0/0</sup> mice. This suggests that the chimeric transgene or some modified version may be superior to HuPrP in terms of generating mice with the shortest incubation times for bioassays of Hu prions.

#### Transmission Of Chimeric Prions

Species specific amino acid variations in PrP are known to contribute significantly to the "species barrier" Prusiner, S.B., et al., "Transgenetic studies implicate interactions between homologous PrP isoforms in scrapie prion replication," Cell 63:673-686 (1990); Scott, M., Foster, D., Mirenda, C., Serban D., Coufal, F., Wälchli, M., Growth, D., Carlson, G.,

- 81 -

DeArmond, S.J., Westaway, D., and Prusiner, S.B. (1989). Transgenic mice expressing hamster prion protein produce species-specific infectivity and amyloid plaques. Cell 59:847-857. Prolongation of incubation times on primary  
5 passage of prions between species is generally seen while second passage in the same species results in a shortening and stabilization of incubation times  
Pattison, I.H., "Experiments with scrapie with special reference to the nature of the agent and the pathology of  
10 the disease," Slow, Latent and Temperate Virus Infections, NINDB Monograph 2, D.C. Gajdusek, et al., eds. (Washington, D.C.: U.S. Government Printing), pp. 249-257 (1965). Primary passage of Hu prions from a sporadic CJD case (EC) produced CNS disease in  
15 Tg(MHu2M)5378/FVB with an incubation time of  $218 \pm 5$  d( $\pm$  SEM) (Table 6 Section A). Brains from ill mice were collected and homogenates inoculated into mice from the same Tg line. Passage of these chimeric prions in Tg(MHu2M)5378/FVB mice gave incubation times similar to  
20 those seen with Hu prions on the primary passage (Table 7 Section A). This finding shows that these Tg(MHu2M)5378/FVB mice are completely permissive for Hu prions. Passage of chimeric prions in Tg(MHu2M)5378/Prnp<sup>0/0</sup> mice resulted in a shortening of the  
25 incubation time by ~20% presumably due to the elimination of MoPrP<sup>C</sup>; i.e., ablating the endogenous mouse prion protein gene.

- 82 -

TABLE 7

Serial transmission of chimeric Hu/Mo prions in Tg(MHu2M) mice.

				Incubation Times
				mean d $\pm$ SEM (n/no)
	Recipient Mouse Line	Inoculum <sup>a</sup>	Illness	Death
5	(A) Chimeric prions produced in Tg(MHu2M) mice inoculated with CJD prions			
	Tg(MHu2M) 5378/FVB	MHu2M( $\Delta$ CJD) <sup>b</sup>	220 $\pm$ 3 (7/7) <sup>c</sup>	226 $\pm$ 1(5)
	Non-Tg5378/FVB	MHu2M( $\Delta$ CJD) <sup>b</sup>	>340	
	Tg(MHu2M) 5378/FVB	MHu2M( $\Delta$ CJD) <sup>d</sup>	226 $\pm$ 3 (9/9)	228 $\pm$ 1(6)
10	Non-Tg5378/FVB	MHu2M( $\Delta$ CJD) <sup>d</sup>	>340	
	Tg(MHu2M) 5378/ Prnp <sup>0/0</sup>	MHu2M( $\Delta$ CJD) <sup>d</sup>	189 $\pm$ 4 (8/8)	192 $\pm$ 1(4)
	Tg(MHu2M) 5378/ Prnp <sup>0/0</sup>	MHu2M( $\Delta$ CJD) <sup>d</sup>	183 $\pm$ 5 (7/7)	190 $\pm$ 3(4)
15	(B) Mouse prions produced in Tg(MHu2M) or non-Tg mice inoculated with RML prions			
	Tg(MHu2M) 5378/FVB	Mo (RML)	178 $\pm$ 3 (19/19)	203 $\pm$ 2(14) <sup>e</sup>
	NonTg5378/FVB	Mo (RML)	127 $\pm$ 2 (18/18)	156 $\pm$ 2(5)
	Tg(MHu2M) 5378/FVB	MHu2M(RML) <sup>f</sup>	185 $\pm$ 1 (7/7)	211 $\pm$ 1(3)
20	Tg(MHu2M) 5378/FVB	MHu2M(RML) <sup>g</sup>	189 $\pm$ 2 (7/7)	211 $\pm$ 9(3)
	Non-Tg5378/FVB	MHu2M(RML) <sup>g</sup>	134 $\pm$ 3 (5/5)	N.D.
	Tg(MHu2M) 5378/ Prnp <sup>0/0</sup>	Mo (RML)	>340	
25	Tg(MHu2M) 5378/ Prnp <sup>0/0</sup>	MHu2M(RML) <sup>f</sup>	>300	
	Tg(MHu2M) 5378/ Prnp <sup>0/0</sup>	MHu2M(RML) <sup>g</sup>	>300	

30 <sup>a</sup> Notation in parentheses indicate inoculum used in initial passage in Tg(MHu2M) mice.

- 83 -

- ° Mice were inoculated with chimeric prions generated in the brain of a Tg(MHu2M)5378/FvB mouse that had been inoculated with a brain homogenate prepared from patient EC who died of sporadic CJD.
- 5 ° Number of mice developing CNS illness divided by the number inoculated are given in parentheses.
- ° Mice were inoculated with chimeric prions generated in the brain of a second Tg(MHu2M)5378/FVB mouse that had been inoculated with a brain homogenate prepared from patient EC who died of sporadic CJD.
- 10 ° Data from (Telling et al. 1994).
- ° Mice were inoculated with Mo prions generated in the brain of a Tg(MHu2M)5378/FVB mouse that had been inoculated with RML Mo prions.
- 15 ° Mice were inoculated with Mo prions generated in the brain of a second Tg(MHu2M)5378/FVB mouse that had been inoculated with RML Mo prions.

#### Specificity Of Chimeric Prions And Transgenes

Non-Tg5378/FVB littermates, which express only MoPrP<sup>C</sup>, inoculated with the chimeric prions have remained well for >340 days. Thus it appears that homology between the substrate PrP<sup>C</sup> and the product PrP<sup>Sc</sup> in the region bounded by residues 96 to 167 is essential for prion propagation. Conversely, Tg(MHu2M)Prnp<sup>0/0</sup> mice are resistant to Mo prions; they have remained well for >340 days after inoculation (Table 7 Section B).

Although Tg(MHu2M)5378/FVB mice are permissive for Mo(RML) prions, the incubation time of  $178 \pm 3$  d( $\pm$  SEM) was protracted compared to that of  $127 \pm 2$  d( $\pm$  SEM) for non-Tg5378/FVB littermates (Table 7 Section B). Two homogenates derived from Tg(MHu2M)5378/FVB mice were inoculated with Mo(RML) prions were passaged in Tg(MHu2M)5378/FVB mice and non-Tg littermates. The incubation time in the Tg(MHu2M)5378/FVB mice did not change while the incubation time in the non-Tg mice

- 84 -

shortened to the incubation time registered for primary passage of Mo(RML) prions in these mice (Table 7 Section B). This behavior and the fact that MoPrP<sup>Sc</sup> is made in response to inoculation with Mo prions (Telling et al., 1994) appears to show that Tg(MHu2M)5378/FVB mice propagate Mo prions from endogenous MoPrP<sup>Sc</sup> and not from MHu2MPrP<sup>Sc</sup>.

Residue 129 Mismatches Between PrP<sup>Sc</sup> In  
The Inoculum And Transgene-encoded PrP<sup>Sc</sup>

- 10 In Caucasians (Palmer et al., 1991) but not Asians Tateishi and Kitamoto, (1993) Developments in diagnosis for prion diseases. Br. Med. Bull. 49,971-979
- 15 homozygosity for M or V codon 129 has been reported to predispose people to development of sporadic CJD.
- 15 Homozygosity at codon 129 in some Baker et al., (1991) Amino acid polymorphism in human prion protein and age at death in inherited prion disease. Lancet 337, 1286; Goldfarb, L.G., et al., "The molecular genetics of human transmissible spongiform encephalopathy", Prion Diseases
- 20 of Humans and Animals, S.B. Prusiner et al., eds. (London: Ellis Horwood), pp. 139-153 (1992) but not other inherited prion diseases diminished the age of onset of CNS dysfunction; Gabizon et al., (1993) Mutation and polymorphism of the prion protein gene in Libyan
- 25 Jews with Creutzfeldt-Jakob disease. Am. J. Hum. Genet. 33, 828-835. The Tg(HuPrP)152 mice express HuPrP with V at codon 129 while another line Tg(HyPrP)440 synthesizes HuPrP with M at 129. When Tg(HuPrP)152/Prnp<sup>0/0</sup> and Tg(HuPrP)440/Prnp<sup>0/0</sup> mice were inoculated with prions from
- 30 iatrogenic and sporadic cases, the shortest incubation times occurred when the amino acid residues at position 129 were the same in PrP<sup>Sc</sup> and inoculated PrP<sup>Sc</sup>. Tg(HuPrP)440/Prnp<sup>0/0</sup> mice inoculated with a case of iatrogenic CJD from a patient with a M/M codon 129



- 85 -

haplotype, referred to as 364, exhibited a mean incubation time of  $164 \pm 2$  d ( $\pm$  SEM) while the same inoculum produced disease in Tg(HuPrP)152/Prnp<sup>0/0</sup> mice with a mean incubation time of  $253 \pm 6$  d ( $\pm$  SEM). Two cases of sporadic CJD derived from patients with the M/M codon 129 haplotype, referred to as EC and RG, produced disease in Tg(HuPrP)152/Prnp<sup>0/0</sup> mice with mean incubation times of  $254 \pm 2$  d ( $\pm$  SEM), respectively (Table 5 Section B).

#### 10 Tg(MHu2M-P101L) Mice Expressing The GSS Mutation

To produce a model of GSS, we created lines of mice carrying the P102L point mutation in both the MoPrP and HuPrP genes. The Tg(MoPrP-P101L)87 and 174 mice expressing the mutant PrP<sup>C</sup> at high levels developed disease spontaneously between 50 and 300 d of age (Hsiao et al., 1994; Hsiao et al., 1990). In contrast, a line designated Tg(HuPrP-P102L)/FVB was observed for >700 d and unlike the Tg(MoPrP-P101L) mice, did not develop spontaneous neurologic disease.

20 The successful transmission of Hu prions to Tg(MHu2M)5378/FVB mice promoted us to produce Tg(MHu2M-P101L)69/Prnp<sup>0/0</sup> mice. Unlike the Tg(HuPrP-P102L) mice, these Tg(MHu2M-P101L) mice spontaneously developed neurologic disease. The mean age of illness in Tg(MHu2M-P1201L) mice was  $362 \pm 13$  d ( $\pm$  SEM). By 480 days, -90% of Tg(MHu2M-P1201L) mice developed CNS dysfunction (n/n<sub>0</sub> = 15/17). An intense reactive astrocytic gliosis was found in the gray matter of all mice expressing the MHu2M-P101L transgene at the time they exhibited signs of CNS dysfunction. Modest spongiform degeneration and PrP immunoreactivity were found in the white matter of all mice examined. Besides the Tg(HuPrP-P102L)7/FVB mice, additional controls include Tg(HuPrP)/FVB, Tg(MHu2M)/FVB

- 86 -

and Tg(MHu2M)/Prnp<sup>0/0</sup> mice, none of which have developed CNS degeneration spontaneously.

Transmission Of GSS Human Prions To Tg(MHu2M-P101L) Mice

Although Tg(MHu2M-P101L)69Prnp<sup>0/0</sup> mice eventually  
5 develop a spontaneous neurologic disorder, we attempted to determine whether the illness would appear more rapidly if the animals are inoculated. Both wt and GSS Hu prions were inoculated. Tg(MHu2M-P101L)69Prnp<sup>0/0</sup> mice were inoculated at about 70 days of age with brain  
10 extract from a GSS patient referred to as JJ, who carried the P102L mutation, or with brain extracts from two sporadic CJD cases (RG and EC in Table 5). These mice inoculated with prions from the GSS patient JJ died after 171 ± 2.8 d (± SEM). The mean age of 247 ± 3 d (± SEM) at  
15 which these Tg mice became ill was more than 100 days earlier than the age at which uninoculated controls developed signs of CNS dysfunction. Although the Tg(MHu2M-P101L) mice inoculated with prions from the sporadic CJD cases have a mean incubation time of 259 ±  
20 10 d (± SEM) (n/n<sub>0</sub> = 12/15), these mice were 350 ± 11 d ± SEM) of age at the time of death. The age of these mice prevented us from concluding whether they became ill from the inoculated prions or spontaneously as a result of the MHu2MPrP-P102L mutant protein.

25 Our findings demonstrate that Hu prions from the GSS patient carrying the point mutation homologous to that in the transgene caused disease more rapidly than did wt Hu prions from sporadic cases of CJD. Conversely, the Hu prions from the GSS patient have failed to produce  
30 disease >280 d after inoculation in Tg(MHu2M)5376/Prnp<sup>0/0</sup> mice (Table 6 Section C); whereas, Hu prions containing wt PrP<sup>Sc</sup> cause disease in Tg(MHu2M)5378/Prnp<sup>0/0</sup> mice at ~190 d (Table 6 Section B). The onset of illness in the

- 87 -

GSS inoculated mice was relatively synchronous, with a range of 30 d while the onset was less uniform in the spontaneously ill and CJD-inoculated Tg(MHu2M-P101L)69/Prnp<sup>0/0</sup> mice with ranges of 210 d and 157 d, respectively.

Tg(MHu2M-P101L) mice inoculated with GSS prions exhibited spongiform degeneration and reactive astrocytic gliosis similar to uninoculated Tg(MHu2M-P101L) mice that developed CNS dysfunction spontaneously. However, the inoculated mice showed more neuronal loss and more intense and widespread GFAP immunostaining than uninoculated, spontaneously ill mice. PrP accumulation was more intense in some gray matter regions such as the hippocampus in the Tg(MHu2M-P101L) mice inoculated with GSS prions than the uninoculated animals exhibiting spontaneous illness.

Uninoculated Tg(MHu2M-P101L)69/Prnp<sup>0/0</sup> mice that developed spontaneously did not have any detectable protease-resistant PrP (PrP 27-30) on Western blots. This finding is similar to that reported previously with Tg(MoPrp-P101L)87 and 174 mice that also develops CNS illness spontaneously Hsiao, K.K., Groth, D., Scott, M., Yang, S.-L., Serban, H., Rapp, D., Foster, D., Torchia, M., DeArmond S.J., and Prusiner, S.B. (1994). Serial transmission in rodents of neurologic disease from transgenic mice expressing mutant prion protein. Likewise, the brain of the GSS patient JJ from which the inoculum was derived contained relatively little or no detectable PrP 27-30 even though numerous PrP amyloid plaques were found Hsiao, K., Baker, H.F., Crow, T.J., Poulter, M., Owen, F., Terwilliger, J.D., Westaway, D., Ott, J., and Prusiner, S.B. (1989). Linkage of a prion protein missense variant to Gerstmann-Sträussler syndrome. Nature 338:342-345. On some occasions, a weak, diffuse band comigrating with PrP 27-30 has been

- 88 -

observed with homogenates prepared from the brain of patient JJ. Whether regional variations in protease-resistant PrP are responsible for these inconsistent results remains to be established. In addition, no PrP 27-30 was found in the brains of the Tg(MHu2M-P101L)69/Prnp<sup>0/0</sup> mice inoculated with homogenate prepared from the brain of the GSS patient JJ at the time they were sacrificed after development of CNS dysfunction. The relatively short incubation times in the Tg(MHu2M-P101L)69/Prnp<sup>0/0</sup> mice argue that the brain of JJ contained high prion titers even if PrP 27-30 was difficult to detect. From these results, we conclude that PrP<sup>Sc</sup> containing the P102L mutation is probably less protease-resistant than wtPrP or PrP carrying other mutations.

15 Transmission of Familial CJD  
(E200K) Human Prions To Tg(MHu2M) Mice

Brain extracts were prepared from two patients who carried the E200K mutation and died of CJD (Gabizon et al., 1993). The extracts were inoculated into Tg(MHu2M)5378/Prnp<sup>0/0</sup> mice that developed CNS dysfunction in 170 ± 2 d (± SEM, n = 10) for the LJ1 case and ~160 d for the CA case. In contrast to the P102L mutation, Hu prions from patients who carried the E200K mutation caused disease as rapidly in Tg(MHu2M)5378/Prnp<sup>0/0</sup> mice as Hu prions containing wtPrP<sup>Sc</sup> from sporadic CJD cases (Table 6 Section C).

Transgenics - Alternative Methods of Producing

Transgenic mice expressing moderate to high levels of wild-type human prion (HuPrP) were originally constructed by microinjecting fertilized FVB embryos with cosmid DNA expressing human PrP. The results of a large number of transmission experiments with sporadic, iatrogenic and familial prion cases revealed that these mice were no more responsive to human prions than their

- 89 -

non-transgenic counterparts. We have demonstrated that by eliminating endogenous mouse (Mo)PrP expression in these transgenic mice, transmission of human prions becomes efficient with mean incubation times as low as 5 160 days. Expression of even half the normal amount of mouse PrP was sufficient to inhibit human prion propagation. These results demonstrate that Mo PrP is extremely inhibitory for the propagation of human prions in transgenic mice even though the level of expression of 10 HuPrP was approximately 8 to 16-fold higher than Mo PrP. These and the results of other genetic experiments have led to the notion that a third component, which we refer to as protein X, must feature in prion propagation. Evidence points to the C-terminal region of PrP as the 15 location for the protein X binding site.

The results of these experiments demonstrate that current transgenic mouse models for the assay of human prions can be improved upon substantially. Because of the inhibitory effects of MoPrP in mice expressing 20 heterologous transgenes, eliminating its expression is crucial for the efficient propagation of heterologous transgenes, eliminating its expressing is crucial for the efficient propagation of heterologous prions in these transgenic mice. This can be achieved in one of several 25 ways.

Homologous Recombination -  
Producing Transgenic Mice  
Crossed With MoPrP Gene Ablated Mice

FVB mice expressing human PrP genes have been 30 constructed using the cos.SHaTet cosmid expression vector derived from the Syrian hamster (SHa). The FVB strain of mice contain and express the normal complement of MoPrP genes and so one method for introducing the HuPrP transgene array into a background in which MoPrP

- 90 -

expression is ablated is by genetic crosses between the transgenic FVB-derived line and a second line of transgenic mice in which both MoPrP genes were disrupted. Mice homozygous for the disrupted Prnp genes were

5 created. These genetically-altered mice were created by a process known as homologous recombination (Thomas and Capecchi, Cell 51:503-512, 1987) in which a selectable disrupted MoPrP gene was introduced into embryonic stem (ES) cells from SV129 mice. Blastocysts of C57BL/6J mice

10 were injected with SV129 ES cells in which one copy of the MoPrP gene had been disrupted thus generating a chimeric mouse with one disrupted allele. That mouse was mated with a C57BL mouse and the offspring crossed to each other to produce null animals in which both copies

15 of the MoPrP gene were disrupted, referred to as Prnp<sup>0/0</sup> mice. Subsequently, these Prnp<sup>0/0</sup> mice were repeatedly crossed onto the FVB background. FVB-derived transgenic mouse lines Tg(HuPrP)FVB/152 and Tg(HuPrP)FVB/440 were crossed with Prnp<sup>0/0</sup> mice. Backcrossing these mice

20 produced animals in which the only PrP<sup>c</sup> molecules that were synthesized were those encoded by the transgene.

Producing Transgenic Mice Using  
Fertilized Oocytes From MoPrP Gene Ablated Mice

The second method for producing transgenic mice in

25 which the only PrP<sup>c</sup> molecules synthesized are encoded by the HuPrP transgene is by directly microinjecting DNA from a vector capable of directing expression of HuPrP. Derivatives of the cos.SHaTet cosmid expression vector containing the HuPrP open reading frame were used --

30 (other expression systems could be used including a cosmid consisting of the cognate HuPrP gene or other vectors capable of appropriate expression of HuPrP in transgenic mice). Using embryos from the originally created C57BL-derived Prnp<sup>0/0</sup> mice we encountered great



- 91 -

difficulty in producing transgenic mice by this method because of the poor survival rates of microinjected embryos. These  $\text{Prnp}^{0/0}$  mice were subsequently repeatedly crossed onto the FVB background to produce mice which  
5 were genetically ~95% FVB but which were also homozygous for the gene ablation. By modifying the  $\text{Prnp}^{0/0}$  mice in this way we now have very high rates of production of transgenic mice by this method.

#### Gene Replacement

- 10 A different approach to eliminating the inhibitory effects of MoPrP would be to create new lines of transgenic mice in which the endogenous MoPrP genes were replaced with HuPrP genes by homologous recombination in ES cells. This gene-replacement approach (Hasty et al.,  
15 Nature 350:243-6, 1991; Valancius and Smithies, Mol. Cell Biol. 11:1402-8, 1991) is a variation of the gene-insertion experiment described above in which  $\text{Prnp}^{0/0}$  mice were created. In gene replacement, the sequences in the input DNA completely replace those in the target DNA.
- 20 The methodologies that are currently available permit gene targeting at high efficiency and fidelity so that it should in theory be possible to replace the MoPrP gene with the homologous HuPrP gene in ES stem cells and thereby produce mice that are homozygous for this  
25 replacement.

After completing gene replacement with either HuPrP or chimeric MHu2MPrP, these mice are mated to transgenic mice expressing high levels of the homologous protein such as HuPrP or MHu2MPrP. The mice will express the  
30 highest levels of the foreign PrP of interest and possess the shortest incubation times. For example, mice with ~50 copies of the MoPrP transgene have incubation times of ~60 days after inoculation with  $\sim 10^6$  ID<sub>50</sub> units if the endogenous MoPrP genes are ablated; in contrast,

- 92 -

incubation times of ~48 days were found if the endogenous MoPrP genes are left intact (Example 8). Alternatively, the fertilized eggs from these mice with gene replacements can be microinjected with the DNA encoding either the same PrP gene as that replaced or a related gene.

#### Creating Mammals Which Express Protein X

Identifying the amino acid sequence(s) in MoPrP responsible for the binding of mouse protein X and construct a modified MoPrP gene in which the sequence for this binding site is mutated. Such a benign MoPrP molecule will not interfere with human prion propagation in transgenic mice expressing HuPrP because protein X is not sequestered by the mutant MoPrP. Following the above-described procedures in which the MoPrP gene is replaced with HuPrP or modified MoPrP genes, it is possible to create transgenic mice expressing HuPrP using these new genetic backgrounds either by genetic crosses or by direct microinjection of a vector capable of directing expression of HuPrP into fertilized embryos from these newly-created transgenic mouse lines.

#### Use Of Multiple Promoters To Increase The Level Of PrP Gene Expression

Further modifications of the current transgenic mouse models for the assay of human prions involve the production of transgenic mice expressing HuPrP under the control of promoter/enhancer sequences from genes other than PrP which normally express their gene products in the central nervous system (CNS). Examples of such promoters include neuron specific enolase (NSE) (Farlie et al., Natl. Acad. Sci USA 92:4397-4401, 1995) or platelet-derived growth factor (PDGF) (Sasahara et al., Cell 64:217-227, 1991). Expression of HuPrP under these

- 93 -

circumstances will result in levels of expression much higher than normally achieved by PrP promoter/enhancer sequences leading to greatly shortened human prion incubation times in transgenic mice expressing these  
5 constructs. Alternatively, by creating several lines of transgenic mice in which HuPrP expression is directed by a variety of different gene constructs and then mating these mice to produce lines in which HuPrP was expressed simultaneously in a variety of different cell types in  
10 the CNS, lines of mice may be created in which HuPrP expression levels are extremely high and incubation times are exceedingly short.

Mice Expressing Multiple PrP Different  
Transgenes To Increase The Range  
15 Of Prions To Which They Are Susceptible

Create lines of transgenic mice simultaneously expressing several different forms of HuPrP derived from patients with polymorphisms and/or with mutations at particular amino acid residues of HuPrP. We have  
20 recently discovered that human prion propagation, at least in the case of the polymorphism at codon 129 and the GSS mutation at codon 102, occurs most efficiently when the amino acid sequences at these residues are matched in PrP<sup>Sc</sup> in the inoculum and PrP<sup>C</sup> encoded by the  
25 transgene. By creating a line in which a number of different forms of HuPrP are expressed, prions with a variety of different polymorphisms and/or mutations are transmitted efficiently to the same host.

- 94 -

TABLE 8

Prion incubation times in Tg(MoPrP-A) mice

				<u>Incubation Times</u>
				mean d $\pm$ SEM (n/no)
	<u>Recipient Mouse</u>	<u>Inoculum<sup>a</sup></u>	<u>Illness<sup>b</sup></u>	<u>Death<sup>c</sup></u>
	(A) Tg(PrP-A) mice			
5	Tg(MoPrP-A) 4053	None	710 $\pm$ 37 (4/20)	N.D.
	Tg(MoPrP-A) 4053	RML	48 $\pm$ 1 (21/21)	53 $\pm$ 2 (12)
	Tg(MoPrP-A) 4053	Tg4053-2 (RML)	48 $\pm$ 2 (10/10)	58 $\pm$ 2 (7)
	Non-Tg 4053	RML	127 $\pm$ 2 (18/18)	156 $\pm$ 2 (5)
10	Tg(MoPrP-A) 4053/ Prnp <sup>0/0</sup>	RML	59 $\pm$ 2 (9/9)	65 $\pm$ 2 (6)
	(B) Swiss CD-1 mice			
	Swiss CD-1 <sup>e</sup>	RML	138 $\pm$ 1 (24/24)	159 $\pm$ 3 (24)
	Swiss CD-1	Tg4053-1 (RML)	140 $\pm$ 3 (10/10)	163 $\pm$ 2 (6)
	Swiss CD-1	Tg4053-2 (RML)	138 $\pm$ 5 (9/9)	169 $\pm$ 2 (6)

15

a Notation in parentheses indicate inoculum used in initial passage. Single digit following a hyphen distinguishes individual mice from a particular line.

20 b The number of mice developing clinical signs of scrapie is shown in parentheses.

25 c The number of mice dying of scrapie is shown in parentheses. Mice that were sacrificed for pathologic examination were excluded from these calculations.

d Data from Butler, et al., J. Virol. 62:1558-1584, 1988.

Further Examples Summarized In Table 9

Above we have described using homogenized brain tissue from human patients dying of human prion disease, either sporadic or iatrogenic Creutzfeld-Jacob Disease (CJD). By inoculating mice of the invention we have shown that it is possible to transmit human CJD to mice -- specifically mice containing the chimeric gene MHu2MPrP. This chimeric gene is a mouse PrP gene with the codons located between codon 94 and codon 188 replaced with corresponding codons from a human PrP gene. The chimeric PrP gene MHu2M differs from the endogenous mouse PrP gene by nine amino acids in the central location which amino acids are, of course, derived from the human PrP gene. Transgenic mice were created using the chimeric gene and using microinjection of fertilized mouse embryos from an inbred mouse line called FVB. The resulting transgenic mice are referred to as Tg(MHu2M)5378/FVB or 5378 mice. The brains of these transgenic mice, in addition to expressing endogenous mouse PrP also expressed transgene-derived MHu2MPrP at levels equivalent to PrP<sup>C</sup> expressed in human brain.

The transgenic mice Tg(MHu2M)5378/FVB mice were then crossed with a line of mice having their endogenous PrP gene ablated. Specifically, they were crossed with Prnp-Abl mice to create a line of transgenic hybrid mice which only express the MHu2MPrP gene. These hybrid transgenic mice are referred to as Tg(MHu2M)5378/Abl. The expression of the MHu2MPrP transgene in the ablated endogenous PrP background provides shorter incubation times when the mice were inoculated with human prions due to the elimination of the interfering effects of endogenous mouse PrP expression. Attempts to breed these 5378/Abl lines to obtain mice homogenous for the transgene were not successful in that the mice developed an unrelated neurologic disorder. Such is not believed

- 96 -

to be due to high level expression of the PrP transgene in that other mice expressing high levels of the transgene did not show symptoms of this disorder.

Table 9 provides a summary of the transmission  
5 experiments into Tg(MHu2M)5378 FVB and the hybrid mice  
Tg(MHu2M)5378/Abl. The results with respect to the  
transgenic 5378 mice appear in column 3 and the results  
for the hybrid transgenic (5378/Abl) mice appear in  
column 4. A shortened incubation time is demonstrated by  
10 the hybrid mice due to the ablated endogenous PrP gene.

A control was run and the transgenic mice were  
inoculated with homogenized brain material from a human  
that did not die from prion related disease but rather  
from amyotrophic lateral sclerosis (ALS). Mice  
15 inoculated with this material did not show symptoms after  
530 days.



TABLE 9

Inoculum	Patient Genotype	TG (MHu2M) 5378/FVB	Tg (MHu2M) 5378/Abl
RG (sCJD)	M/M, 129	238 $\pm$ 3 (8/8)	191 $\pm$ 3 (10/10)
5 EC (sCJD)	M/M, 129	218 $\pm$ 5 (7/7)	
364 (iCJD)	M/M, 129	232 $\pm$ 3 (9/9)	191 $\pm$ 6 (8/8)
MA (sCJD)	M/M, 129	222 $\pm$ 1 (4/4)	180 $\pm$ 5 (8/8)
RC (sCJD)	M/M, 129		207 $\pm$ 4 (8/8)
WL (sCJD)	M/M, 129		181 $\pm$ 5 (7/10)
10 MHuM-1 (EC, sCJD)		220 $\pm$ 3 (7/7)	189 $\pm$ 4 (8/8)
MHuM-2 (EC, sCJD)		226 $\pm$ 3 (9/9)	183 $\pm$ 5 (7/7)
RO (sCJD)	M/M, 129		217 $\pm$ 2 (9/9)
E200K-1 (fCJD)	M/M, 129 and E200K		179 $\pm$ 2 (10/10)
E200K-2 (fCJD)	M/M, 129 and E200K		179 $\pm$ 1 (8/8)
15 E200K-3 (fCJD)	M/M, 129 and E200K		184 $\pm$ 4 (8/8)
E200K-4 (fCJD)	M/M, 129 and E200K		180 $\pm$ 9 (9/9)
E200K-5 (fCJD)	M/V, 129 and E200K		>650 days (0/4)

Experiments re Results Shown in Table 10

Hybrid mice of the type described above were  
 20 produced. Specifically, the hybrid mice with an ablated  
 endogenous PrP gene which were then further modified by  
 the addition of the chimeric mouse/human PrP gene.  
 However, these mice were produced in a manner so as to  
 obtain a high copy number of the chimeric mouse/human PrP  
 25 gene in an attempt to reduce incubation times. The  
 resulting mice expressed approximately 8 to 16 fold  
 higher levels of the chimeric gene. However, the mice

- 98 -

showed longer incubation times as compared with the original hybrid mice with lower copy numbers of the chimeric gene. The results are shown below in Table 10.

TABLE 10

5	Inoculum	Patient Genotype	Tg (MHu2M) 3095/Abl	Tg (MHu2M) 3096/Abl	Tg (MHu2M) 3305/Abl
	RG (sCJD)	M/M,129	239 $\pm$ 4 (11/11)		
	EC (sCJD)	M/M,129		237 $\pm$ 5 (11/11)	
	364 (iCJD)	M/M,129			271 $\pm$ 10 (8/8)
	MA (sCJD)	M/M,129	264 $\pm$ 7 (12/12)		
10	RC (sCJD)	M/M,129		266 $\pm$ 8 (11/11)	
	RO (sCJD)	M/M,129			234 $\pm$ 5 (10/10)

#### Additional Mouse/Human PrP Chimeric Genes

In order to more closely focus on the particular codons within the PrP gene which are related to the development of prion disease additional chimeric human/mouse PrP genes were created. The three new chimeric constructs are referred to as MHu3, Hu3M and Hu4M PrP. In each of the new constructs a different part of the mouse PrP gene has been replaced with corresponding codons from a human PrP gene. When these chimeric genes are inserted within the mice such will provide further information with respect to which sequences facilitate the conversion of PrP<sup>c</sup> to PrP<sup>sc</sup> more efficiently and therefore result in shorter incubation times. It is possible to change the codons and thus the

- 99 -

amino acid residues throughout the genes. However, attention is focused on the central region of the gene. Further, it is possible to change either single amino acids or entire segments.

5 Results re Table 11 (Mutation at PrP Codon 129)

The most common form of inherited human prion disease involves a mutation at codon 200 of the PrP gene. This mutation results in the replacement of the amino acid glutamate with lysine. The mutation is found most  
10 frequently among the population of Libyan Jews residing in Israel. Families having this mutation have been observed in other parts of the world. Homogenized brain tissue from four different individuals dying with the disease have been used as an inoculant in the above-  
15 described hybrid transgenic mice (specifically Tg(MHu2M)5378/Abl mice). The incubation times range between 160 and 180 days. However, when an inoculant was used from a fifth individual who was a Libyan Jew with the mutation at codon 200 the mice did not demonstrate  
20 symptoms of disease after 620 days.

The ability to transmit the disease from the first four individuals but not from the fifth is believed to be related to differences in the PrP gene between the individuals at codon 129. Specifically, the four  
25 individuals whose brain tissue did allow for the transmission of disease to the hybrid mice were all homozygous for methionine (MET/MET) at codon 129. However, the fifth individual whose brain tissue was not able to transmit symptoms into the hybrid mice was  
30 heterozygous Met\Val at codon 129.

Speculation about the effect of codon 129 was confirmed by further experimentation wherein the 5378/Abl hybrid mice were inoculated with homogenized brain tissue from four different humans who died of fatal familial  
35 insomnia (FFI) and a case of familial CJD. Transmission

- 100 -

was observed to mice from three of the individuals with FFI but not the fourth individual with FFI. The FFI disease includes a mutation of the human PrP gene at codon 178 resulting in the replacement of the amino acid aspartate with asparagine. The FFI disease is seen in individuals when the mutated allele encodes methionine at codon 129. The same mutation causes a form of familial CJD, but only when the mutated allele encodes valine at codon 129. Inoculation of the 5378/Abl hybrid transgenic mice with familial CJD with codon 178 mutation has not resulted in the demonstration of symptoms of prion disease after 280 days. These experiments further emphasize the importance of codon 129 with respect to the transmissibility of human prion disease.

Brain tissue was extracted from two human individuals who died of GSS which individuals showed a mutation of the human PrP gene at codon 102 which mutation resulted in replacement of Proline with Leucine. The brain tissue was used to inoculate the 5378/Abl hybrid transgenic mice described above and the mice did not show symptoms of disease after 560 days.

Different transgenic hybrid mice were then produced which mice were similar to the 5378/Abl mice but which mice had their PrP gene changed so that the PrP gene encoded Leucine instead of Proline at codon 101. When these mice were inoculated with the homogenized brain tissue from the same individuals who died of GSS the mice showed symptoms of disease at approximately 200 days as shown in Table 11 below. These experiments show that homology between PrP<sup>Sc</sup> in the inocula and the transgene-expressed PrP appears to be necessary for a efficient transmission of human prions to the transgenic mice. Specifically, the mice Tg(MHu2M-P101L)69/Abl did demonstrate symptoms apparently due to the mutation at

- 101 -

codon 101 which matched the mutation of the humans who died of prion disease.

Brain tissue from other individuals dying of GSS were obtained wherein those individuals showed mutations  
5 at codons 117, 145 and 217. This material was used to inoculate the 5378/Abl mice and did not result in producing symptoms. Mice could be produced wherein the PrP gene of the transgenic hybrid mice had mutations matching the human individuals which died (at codons 117,  
10 145 and 217) and such mice would be expected to show symptoms of disease after approximately 200 days ( $\pm$  50 days) being inoculated with homogenized brain tissue from the GSS individuals.

TABLE 11

	Inoculum	Patient Genotype	Tg (HuPrPV129) 152HOZ/Abl	Tg (MHu2M) 5378/Abl
	RG (sCJD)	M/M, 129	212 ± 3 (8/13)	
	EC (sCJD)	M/M, 129	204 ± 5 (4/14)	
5	RP (sCJD)	V/V, 129	204 ± 6 (4/12)	
	D178N (FFI) Patient A, Region 1	M/M, 129 and D178N		193 ± 5 (9/9)
10	D178N (FFI) Patient B, Region 1	M/V, 129 and D178N		246 ± 0 (1/10)
	D178N (FFI) Patient B, Region 8	M/V, 129 and D178N		>250 ± 0 (0/10)
15	D178N-C (FFI) Patient C, Region 1	M/V, 129 and D178N		222 ± 7 (7/9)
20	D178N (fCJD) Patient D, Region 6	M/M, 129 and D178N		206 ± 7 (7/7)
	D178-D (FFI) Patient D, Region 7	M/M, 129 and D178N		213 ± 0 (1/10)
25	D178N-D (FFI) Patient D, Region 1	M/M, 129 and D178N		232 ± 9 (5/7)
	D178N (CJD)	V/V, 129 and D178N		>280 days 0/10
	JJ (GSS)	V/V, 129 and P102L		>400 days (0/10)
30	JJ (GSS)	V/V, 129 and P102L		>250 days (0/10)
	JB (GSS)	M/M, 129 and P102L		>250 days (0/8)
	GSS	V/V, 129 and A117V		>250 days (0/9)
	GSS	V/V, 129 and Q217R		>560 days (0/8)



- 103 -

Results re Table 12 (Further Mutations)

Based on the results shown above in Table 11 it can be concluded that certain genetic manipulations in the PrP gene of a transgenic mouse are desirable when using the transgenic mice to detect the presence of prions from a specific infected individual. More specifically, it can be seen that it is desirable to construct a chimeric gene of a transgenic mouse so as to include a mutation at a particular point (e.g., at codon 101 of a mouse PrP gene which corresponds to codon 102 of a human PrP gene) which point mutation matches a natural mutation in an individual who died of prion infection and whose brain tissue is used as inoculant in the transgenic mouse. By matching the mutation in the PrP chimeric gene of the transgenic mouse with the mutation in the individual from whom the inoculant is obtained it is possible to detect the presence of prions in the inoculant. Thus, a transgenic mouse with several different PrP genes (each with a different codon mutation) or a gene with several different point (codon) mutations would be expected to develop disease when inoculated with an inoculant containing all or any one of the prions from individuals with different genetic mutations in the PrP gene.

TABLE 12

Inoculum	Patient Genotype	Tg (MHu2M) 5378/Abl	Tg (MHu2M-P101L) 69Abl
JJ(GSS)	V/V, 129	>400 days (0/10)	171 $\pm$ 3 (10/10)
JJ(GSS)	V/V, 129	>250 days (0/10)	164 $\pm$ 10 (9/9)
JB	M/M, 129	>250 days (0/10)	170 $\pm$ 5 (15/15)

- 104 -

Results re Table 13

Transgenic mice expressing a human PrP gene were originally produced using embryos of FVB mice via microinjection of DNA. Two different lines of mice expressing human PrP gene with valine at codon 129 were used for transmission experiments. The lines are referred to as Tg(HuPrP, V129)152/FVB and 110/FVB. In general, the success rate in transmission of human prions to these transgenic mice was found to be no higher than transmission of human prions to non-transgenic mice. It is pointed out that these mice do not have an ablated endogenous PrP gene and do not include a chimeric mouse/human gene. The results are shown in Table 12. However, an exception was observed for one inoculum wherein the inoculum was obtained from an individual dying of iCJD. This inoculum was capable of transmitting human prions to mice at a rate of about 2 in 10 after 589 days.

TABLE 13

20	Inoculum	Patient Genotype	TG(HuPrP V129) 152/FVB	Tg(HuPrP V129) 110/FVB
	RG(sCJD)	M/M,129	>570 (0/10)	
	364(iCJD)	M/M,129		>800 days (0/10)
	170(iCJD)	M/V,129		589 ± 0 (2/10)
25	D178N-A(FFI)	M/M,129 and D178N	>500 days (0/10)	

Results re Table 14

The transgenic mice described above which included the entire human PrP gene were not susceptible to infection with human prions as shown within Table 13.

- 5 However, these mice were crossed with mice having an ablated PrP gene to produce hybrid mice referred to as Tg(HuPrP, V129)152/Abl mice. The results are shown below in Table 14 indicating that it was possible to transmit human prions to these hybrid mice having the endogenous  
10 PrP gene ablated.

TABLE 14

Inoculum	Patient Genotype	Tg(HuPrP, V129) 152/Abl	Tg(HuPrP, V129) 152HCZ/Ab
RG(sCJD)	M/M,129	263 $\pm$ 2 (6/6)	212 $\pm$ 3 (8/13)
EC(sCJD)	M/M,129	254 $\pm$ 5 (9/9)	203 $\pm$ 5 (4/14)
15 364(iCJD)	M/M,129	262 $\pm$ 8 (6/6)	
RP (sCJD)	V/V,129		204 $\pm$ 6 (4/12)

Summary of Results re Table 15

- The hybrid mice described above with the ablated endogenous PrP gene were mated together to produce  
20 homozygous mice for the transgene array. These mice were referred to as Tg(HuPrP, V129)152 HOZ/Abl. These mice, presumably express twice the level of human PrP. The incubation time for these mice was shorter than the mice described above with respect to Table 13. In addition,  
25 transgenic hybrid mice were produced which expressed methionine 129 from the human PrP gene. These mice were referred to as Tg(HuPrP, M129)440/Abl. These mice demonstrated a very short incubation time when inoculated

- 106 -

with human prions. It would be expected that by breeding these mice to produce homozygous mice would further reduce the incubation time.

TABLE 15

5		Inoculum	Patient Genotype	Tg (HuPrP M129) 440/Abl
		RG (sCJD)	M/M, 129	165 ± 4 (7/7)
		EC (sCJD)	M/M, 129	157 ± 3 (7/7)
		364 (iCJD)	M/V, 129	164 ± 2 (7/7)
10		MHu2M-1 (EC, sCJD)		175 ± 3 (6/6)
		MHu2M-2 (EC, sCJD)		182 ± 2 (9/9)

The instant invention is shown and described herein in what is considered to be a most practical and preferred embodiments. It is recognized, however, that 15 departures may be made from which are within the scope of the invention and that modifications will occur to one who is skilled in the art upon reading this disclosure.

- 107 -

CLAIMS

1. A standardized prion preparation produced by the process comprising:
  - producing a plurality of non-human host mammals each
  - 5 having a genome which is substantially identical in terms of genetic materials related to a PrP gene, which mammals are susceptible to infection with a prion which generally only infect an animal which is genetically diverse from the host animal;
  - 10 inoculating the host mammals with a prion containing composition;
  - observing the mammals until the mammals exhibit symptoms of prion infection;
  - harvesting brain tissue from the mammals exhibiting
  - 15 symptoms of prion infection; and
  - homogenizing the harvested brain tissue to provide a standardized prion preparation.

- 108 -

2. The prion preparation of claim 1, wherein the host mammal is selected from the group consisting of a mouse, a rat, a hamster, a rabbit, and guinea pig;

5 further wherein the exogenous PrP gene comprises a nucleotide sequence corresponding to that of an endogenous PrP gene of a genetically diverse mammal selected from the group consisting of human, cow, sheep and horse;

10 further wherein the host mammals have an artificial PrP gene operatively inserted into their genome which PrP gene renders the host mammal susceptible to infection with prions which normally infect only a genetically diverse test animal, wherein the gene is comprised of a  
15 natural codon sequence of a PrP gene of the host mammal with one or more, but not all, of its codons replaced with a different codon of a natural PrP gene of the genetically diverse test animal;

still further wherein the host mammals have their  
20 endogenous PrP gene ablated and have an exogenous PrP gene operably inserted into their genome.



3. The standardized prion preparation of claim 1, wherein the transgenic host is produced using a chimeric gene, comprising:

C-terminus and N-terminus codon sequences of a PrP  
5 gene of a mammal of a first genus;

codon sequences of a PrP gene of a mammal of a  
second genus operatively connected between the N-terminus  
and C-terminus codon sequences from the mammal of the  
first genus, wherein the chimeric gene renders a mammal  
10 of the first genus susceptible to infection with a prion  
normally infectious only to a mammal of the second genus  
when the chimeric gene is operably inserted into the  
genome of a mammal of the first genus, wherein the first  
genus is selected from the group consisting of Mus,  
15 Rattus, Oryctolagus, and Mesocricetus, and further  
wherein the second genus is selected from the group  
consisting of Bos, Ovis, Sus, and Homo.

4. The standardized prion preparation of claim 3,  
wherein the mammal of the first genus is a mouse and the  
20 mammal of the second genus is a human.

5. A method of testing a sample for the presence  
of prion, comprising:

inoculating a first non-human mammal with the  
sample, wherein the first mammal has an altered genome  
25 rendering the first mammal susceptible to infection with  
a prion which generally only infects a genetically  
diverse mammal;

inoculating a second non-human mammal with the  
standardized prion preparation of claim 1, wherein the  
30 first and second mammals each have a genome which is  
substantially identical in terms of genetic material  
related to prions; and

- 110 -

observing the first and second mammals in order to determine if the mammal develops symptoms of prion infection and using the second mammal as a control to judge when a mammal will develop symptoms of prion  
5 infection when inoculated with prions.

6. The method of claim 5, wherein the sample used to inoculate the first mammal is selected from the group of samples consisting of (a) a pharmaceutical formulation containing a therapeutically active component extracted  
10 from an animal source, (b) a component extracted from a human source, (c) an organ, tissue, body fluid or cells extracted from a human source, (d) a formulation selected from the group consisting of injectables, orals, creams, suppositories, and intrapulmonary delivery formulations,  
15 (e) a cosmetic, and (f) a pharmaceutically active compound extracted from a mammalian cell culture.

7. A transgenic host animal having a genome comprised of a PrP gene rendering the hosts animal susceptible to infection with prions which infect and  
20 cause disease in a genetically diverse test animal;

wherein the PrP transgene is comprised of a PrP gene native to the host animal with one or more of its codons replaced with a corresponding codon of a PrP gene of the genetically diverse test animal and wherein one or more  
25 of the replacing codons of the genetically diverse test animal is a codon indicative of a prion disease in the genetically diverse test animal.

8. The transgenic host animal of claim 7, wherein the host animal is selected from the group consisting of  
30 a mouse, a rat and a guinea pig and the genetically diverse test animal is selected from the group consisting

- 111 -

of a human, a sheep and a cow, and further wherein the endogenous PrP gene of the host animal is ablated.

9. The transgenic host animal of claim 8, wherein the host animal is a mouse and the genetically diverse  
5 test animal is a human and the replacing codon of the human PrP gene is a codon located at a position selected from the group consisting of 102, 129 and 145.

10. The transgenic host animal as claimed in claim 8, wherein the host animal is a mouse, the geentically  
10 diverse test animal is a human and the mouse PrP gene at codon 101 is replaced with a corresponding human codon at position 102.




FIG. 1




FIG. 2

2 / 4

Mo Met Ala Asn Leu Gly Tyr Trp Leu Leu Ala Leu Pro Val Thr Met Trp 1  
 Hu Cys Met Val Ala Thr  
 Mo Thr Asp Val Gly Leu Cys Lys Lys Arg Pro Lys Pro Gly Gly Trp Asn 22  
 Hu Ser Leu  
 Mo Thr Gly Gly Ser Arg Tyr Pro Gly Gln Gly Ser Pro Gly Gly Asn Arg 48  
 Hu  
 Mo Tyr Pro Pro Gln Gly Gly --- Thr Trp Gly Gln Pro His Gly Gly Gly 61  
 Hu Gly Gly  
 Mo Trp Gly Gln Pro His Gly Gly Ser Trp Gly Gln Pro His Gly Gly Ser 79  
 Hu Gly Gly  
 Mo Trp Gly Gln Pro His Gly Gly Gly Trp Gly Gln Gly Gly Gly Thr His 95  
 Hu  
 Mo Asn Gln Trp Asn Lys Pro Ser Lys Pro Lys Thr Asn Leu Lys His Val 111  
 Hu Ser Met Met  
 Mo Ala Gly Ala Ala Ala Ala Gly Ala Val Val Gly Gly Leu Gly Gly Tyr 127  
 Hu  
 Mo Met Leu Gly Ser Ala Met Ser Arg Pro Met Ile His Phe Gly Asn Asp 163  
 Hu Ile Ser  
 Mo Trp Glu Asp Arg Tyr Tyr Arg Glu Asn Met Tyr Arg Tyr Pro Asn Gln 159  
 Hu Tyr His  
 Mo Val Tyr Tyr Arg Pro Val Asp Gln Tyr Ser Asn Gln Asn Asn Phe Val 175  
 Hu Met Glu  
 Mo His Asp Cys Val Asn Ile Thr Ile Lys Gln His Thr Val Thr Thr Thr 191  
 Hu  
 Mo Thr Lys Gly Glu Asn Phe Thr Glu Thr Asp Val Lys Met Met Glu Arg 207  
 Hu  
 Mo Val Val Glu Gln Met Cys Val Thr Gln Tyr Gln Lys Glu Ser Gln Ala 221  
 Hu Ile Glu Arg  
 Mo Tyr Tyr Asp Gly Arg Arg Ser Ser Ser Thr Val Leu Phe Ser Ser Pro 239  
 Hu Gln --- --- Gly Met  
 Mo Pro Val Ile Leu Leu Ile Ser Phe Leu Ile Phe Leu Ile Val Gly 254  
 Hu

Figure 3: Predicted amino acid sequence of mouse PrP and the amino acid differences between mouse and human PrP.

3/4

Mo Met Ala Asn Leu --- --- Gly Tyr Trp Leu Leu Ala Leu Phe Val Thr 14  
 Bo Val Lys Ser His Ile Ser Ile Val Ala  
 Mo Met Trp Thr Asp Val Gly Leu Cys Lys Lys Arg Pro Lys Pro Gly Gly 20  
 Bo Ser  
 Mo --- Trp Asn Thr Gly Gly Ser Arg Tyr Pro Gly Gln Gly Ser Pro Gly 26  
 Bo  
 Mo Gly Asn Arg Tyr Pro Pro Gln Gly Gly --- Thr Trp Gly Gln Pro His 30  
 Bo Gly Gly  
 Mo Gly Gly Gly Trp Gly Gln Pro His Gly Gly Ser Trp Gly Gln Pro His 36  
 Bo Gly  
 Mo Gly Gly Ser Trp Gly Gln Pro His Gly Gly Gly Trp Gly Gln --- --- 40  
 Bo Gly Pro His  
 Mo --- --- --- --- --- --- Gly Gly Gly Thr His Asn Gln Trp Asn Lys 100  
 Bo Gly Gly Gly Gly Trp Gly Gln Gly  
 Mo Pro Ser Lys Pro Lys Thr Asn Leu Lys His Val Ala Gly Ala Ala Ala 116  
 Bo Met  
 Mo Ala Gly Ala Val Val Gly Gly Leu Gly Gly Tyr Met Leu Gly Ser Ala 122  
 Bo  
 Mo Met Ser Arg Pro Met Ile His Phe Gly Asn Asp Trp Glu Asp Arg Tyr 128  
 Bo Leu Ser Tyr  
 Mo Tyr Arg Glu Asn Met Tyr Arg Tyr Pro Asn Gln Val Tyr Tyr Arg Pro 134  
 Bo His  
 Mo Val Asp Gln Tyr Ser Asn Gln Asn Asn Phe Val His Asp Cys Val Asn 140  
 Bo  
 Mo Ile Thr Ile Lys Gln His Thr Val Thr Thr Thr Thr Lys Gly Glu Asn 200  
 Bo Val Glu  
 Mo Phe Thr Glu Thr Asp Val Lys Met Met Glu Arg Val Val Glu Gln Met 212  
 Bo Ile  
 Mo Cys Val Thr Gln Tyr Gln Lys Glu Ser Gln Ala Tyr Tyr Asp Gly Arg 228  
 Bo Gln ---  
 Mo Arg Ser Ser Ser Thr Val Leu Phe Ser Ser Pro Pro Val Ile Leu Leu 244  
 Bo --- Gly Ala Val Ile  
 Mo Ile Ser Phe Leu Ile Phe Leu Ile Val Gly 254  
 Bo

Figure 4: Predicted amino acid sequence of mouse PrP and the amino acid differences between mouse and bovine PrP

REPLACED



Mo Met Ala Asn Leu --- --- Gly Tyr Trp Leu Leu Ala Leu Phe Val Thr 1  
 Sh Val Lys Ser His Ile Ser Ile Val Ala  
 Mo Met Trp Thr Asp Val Gly Leu Cys Lys Lys Arg Pro Lys Pro Gly Gly 11  
 Sh Ser  
 Mo --- Trp Asn Thr Gly Gly Ser Arg Tyr Pro Gly Gln Gly Ser Pro Gly 15  
 Sh  
 Mo Gly Asn Arg Tyr Pro Pro Gln Gly Gly --- Thr Trp Gly Gln Pro His 19  
 Sh Gly Gly  
 Mo Gly Gly Gly Trp Gly Gln Pro His Gly Gly Ser Trp Gly Gln Pro His 23  
 Sh Gly  
 Mo Gly Gly Ser Trp Gly Gln Pro His Gly Gly Gly --- Trp Gly Gln Gly 27  
 Sh Gly  
 Mo Gly Gly Thr His Asn Gln Trp Asn Lys Pro Ser Lys Pro Lys Thr Asn 107  
 Sh Ser --- His Ser  
 Mo Leu Lys His Val Ala Gly Ala Ala Ala Ala Gly Ala Val Val Gly Gly 121  
 Sh Met  
 Mo Leu Gly Gly Tyr Met Leu Gly Ser Ala Met Ser Arg Pro Met Ile His 135  
 Sh Leu  
 Mo Phe Gly Asn Asp Trp Glu Asp Arg Tyr Tyr Arg Glu Asn Met Tyr Arg 155  
 Sh Tyr  
 Mo Tyr Pro Asn Gln Val Tyr Tyr Arg Pro Val Asp Gln Tyr Ser Asn Gln 171  
 Sh  
 Mo Asn Asn Phe Val His Asp Cys Val Asn Ile Thr Ile Lys Gln His Thr 187  
 Sh Val  
 Mo Val Thr Thr Thr Thr Lys Gly Glu Asn Phe Thr Glu Thr Asp Val Lys 201  
 Sh Ile  
 Mo Met Met Glu Arg Val Val Glu Gln Met Cys Val Thr Gln Tyr Gln Lys 219  
 Sh Ile Ile Arg  
 Mo Glu Ser Gln Ala Tyr Tyr Asp Gly Arg Arg Ser Ser Ser Thr Val Leu 233  
 Sh Gln --- --- Gly Ala Val Ile  
 Mo Phe Ser Ser Pro Pro Val Ile Leu Leu Ile Ser Phe Leu Ile Phe Leu 251  
 Sh  
 Mo Ile Val Gly  
 Sh

254

Figure 5: Predicted amino acid sequence of mouse PrP and the amino acid differences between mouse and sheep PrP

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US96/12510

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : A61K 49/00; C12N 7/00, 7/02, 15/00, 15/06

US CL : 424/9.1, 9.2; 435/172.3, 235.1, 239, 320.1; 800/2

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 424/9.1, 9.2; 435/172.3, 235.1, 239, 320.1; 800/2

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Databases: APS, Biosis, CA, Embase, Medline

Search Terms: prion?; prp/ transgen?; homolog?; recomb?; prusiner?/au; scott?/au; telling?/au

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	PRUSINER, S.B. et al. Ablation of the prion protein (PrP) gene in mice prevents scrapie and facilitates production of anti-PrP antibodies. Proceedings of the National Academy of Sciences USA. November 1993, Vol. 90, pages 10608-10612, see entire document.	1-10
Y	SCOTT, M. et al. Propagation of prions with artificial properties in transgenic mice expressing chimeric PrP genes. Cell. 04 June 1993, Vol. 73, pages 979-988, see entire document.	1-10
Y	BUELER, H. et al. Mice devoid of PrP are resistant to scrapie. Cell. 02 July 1993, Vol. 73, pages 1339-1347, see entire document.	1-10

☐ Further documents are listed in the continuation of Box C.☐ See patent family annex.

* Special categories of cited documents	* T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
* A* documents defining the general state of the art which is not considered to be of particular relevance	* X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
* E* earlier document published on or after the international filing date	* Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combinations being obvious to a person skilled in the art
* L* documents which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	* G* documents member of the same patent family
* O* documents referring to an oral disclosure, use, exhibition or other means	
* I* documents published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

15 OCTOBER 1996

Date of mailing of the international search report

28 OCT 1996

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

BRIAN R. STANTON

Telephone No. (703) 308-0196